

Live vaccinations for infants exposed to maternal infliximab in utero and via breast milk – the need for nuanced decision making

Christian Philipp Selinger ¹, Klaartje Bel Kok,² Jimmy K Limdi,^{3,4} Alexandra Kent,⁵ Rachel Cooney ⁶, Catherine Nelson-Piercy ⁷

Infant health outcomes are a key consideration in decisions regarding therapy for pregnant and breast-feeding women with inflammatory bowel disease (IBD). A recent study by Kanis *et al* reported outcomes from 1000 children born to mothers with IBD (196 exposed to anti-TNFs) with no increased risk of adverse outcomes associated with anti-TNF exposure.¹ Anti-TNF therapy should be continued through all trimesters to avoid maternal flares and consequent adverse fetal outcomes.^{2, 3} Discontinuation in the third trimester is not associated with a decrease in infant infection risks.² As the infant will be exposed to anti-TNF by active placental transfer, live vaccinations for the infant should be delayed to avoid potentially fatal infant outcomes.³

The Medicines and Healthcare products Regulatory Agency (MHRA) and European Medicines Agency (EMA) have recently issued guidance for infants exposed to Infliximab in utero and by breast

feeding.⁴ They recommend that infants exposed to Infliximab in utero should not receive live vaccinations until 12 months of age.⁴ As clearance of Infliximab from the infant's blood stream occurs after a mean of 7 months and may take 12 months, this is logical and is reflected in the new European Crohns and Colitis Organisation (ECCO) guidelines.^{3, 5} The second recommendation is that live vaccinations should be avoided in infant exposed to Infliximab through breast milk.⁴ While Infliximab has been detected in breast milk in small amounts, there is no credible evidence that this leads to clinically significant serum levels in infants.⁶ In the few cases where levels were reported in infants, most had in utero exposure. Additionally, even when these drugs are ingested by the baby, they are inactivated by digestive enzymes and poorly absorbed, making exposure less relevant.⁷ In light of the very limited but mainly reassuring evidence we take issue with this blanket recommendation.

Breast feeding has significant benefits from excellent nutrition to potential reductions in childhood infections, and (in women without IBD), a reduced risk of the infant developing IBD.^{1, 8} Despite this, breastfeeding rates are low in anti-TNF exposed mothers.¹ For most women on Infliximab breast feeding their infant, the impact

of the new MHRA and EMA guidance should be limited. Although previous guidance advocated delaying live vaccinations for at least 6 months⁹ and a meta-analysis showed that adverse events from vaccination only occurred when these were given during first 6 months of life,¹⁰ the MHRA recommendation to delay live vaccinations to 12 months aims to maintain safety. This is now supported by the newest ECCO guidelines on reproduction and lactation.³ The main vaccines affected by this are rotavirus (at 8 and 12 weeks) and BCG (4 weeks in high-risk groups) vaccines. However, the first dose of the Measle Mumps Rubella vaccine is given at 12 months.

There are two important scenarios that we feel need careful consideration. First, in cases of in utero exposure and breast feeding for longer than 12 months, should we continue to avoid BCG vaccine in high risk groups? The risk of the infant developing tuberculosis post BCG vaccination associated with breast milk infliximab exposure is probably extremely low. Second, in women without Infliximab exposure during pregnancy, and when Infliximab is started post-partum while breast feeding, the risk would seem similarly low.

In the absence of credible evidence supporting infection risk and adverse events in breast-fed infants of mothers treated with infliximab, the consequences of breast-fed infants not receiving scheduled live vaccines beyond 12 months, or indeed mothers choosing not to breast feed as a result, mandates careful consideration and a nuanced approach. Especially for infants at high-risk of developing tuberculosis if not vaccinated a balanced approach is required as earlier vaccination protect better from acquiring TB through community exposure.

We are surprised that the MHRA and EMA have issued guidance only on Infliximab when similar considerations may be applied to Adalimumab and Golimumab therapy.

¹Gastroenterology, Leeds Teaching Hospitals NHS Trust, Leeds, UK

²Department of Gastroenterology, Barts Health NHS Trust, London, UK

³The Pennine Acute Hospitals NHS Trust Gastroenterology, Manchester, UK

⁴The University of Manchester, Manchester, UK

⁵Gastroenterology, King's College Hospital NHS Foundation Trust, London, UK

⁶Department of Gastroenterology, University Hospitals Birmingham NHS Trust, Birmingham, UK

⁷Guy's and St Thomas' Hospitals NHS Trust, London, UK

Correspondence to Dr Christian Philipp Selinger; christian.selinger@web.de

While Adalimumab may clear earlier from the infant's blood stream after in utero exposure than Infliximab, we feel that the principal safety issues are very similar and clear guidance should be applied to the class of anti-TNF medications used.⁵ The new ECCO guidelines therefore have a simple unified message for all anti-TNF medications and recommend a delay of all live vaccinations for 1 year.³ We caution against a "one size fits all" approach and suggest that, IBD and obstetric teams continue to provide well-considered and evidence based advice on effects of maternal IBD medication on suitability of childhood vaccinations. Clinicians need to weight the risk of vaccination against the risk of acquiring TB through community exposure without discouraging women from breast feeding.

Contributors CPS drafted the commentary and all authors critically reviewed it.

Competing interests 'Yes, there are competing interests for one or more authors and I have provided a Competing Interests statement in my manuscript and in the box below'

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Selinger CP, Bel Kok K, Limdi JK, *et al*. *BMJ Open Gastro* 2022;9:e001029. doi:10.1136/bmjgast-2022-001029

BMJ Open Gastro 2022;9:e001029.
doi:10.1136/bmjgast-2022-001029

ORCID iDs

Christian Philipp Selinger <http://orcid.org/0000-0003-2022-5859>

Rachel Cooney <http://orcid.org/0000-0003-3710-157X>

Catherine Nelson-Piercy <http://orcid.org/0000-0001-9311-1196>

REFERENCES

- Kanis SL, Modderman S, Escher JC, *et al*. Health outcomes of 1000 children born to mothers with inflammatory bowel disease in their first 5 years of life. *Gut* 2021;70:1266–74.
- Luu M, Benzenine E, Doret M, *et al*. Continuous Anti-TNF α use throughout pregnancy: possible complications for the mother but not for the fetus. A retrospective cohort on the French National health insurance database (evasion). *Am J Gastroenterol* 2018;113:1669–77.
- Torres J, Chaparro M, Julsgaard M, *et al*. European Crohn's and Colitis Guidelines on Sexuality, Fertility, Pregnancy, and Lactation. *J Crohns Colitis* 2022.
- Infliximab. (remicade, Flixabi, Inflectra, Remsima and Zessly): use of live vaccines in infants exposed in utero or during breastfeeding. Direct healthcare professional communication (DHPC) medicines and healthcare products regulatory agency, 2022
- Julsgaard M, Christensen LA, Gibson PR, *et al*. Concentrations of Adalimumab and Infliximab in Mothers and Newborns, and Effects on Infection. *Gastroenterology* 2016;151:110–9.
- Fritzsche J, Pilch A, Mury D, *et al*. Infliximab and adalimumab use during breastfeeding. *J Clin Gastroenterol* 2012;46:718–9.
- Namekata T, Miki K, Kimmey M, *et al*. Chronic atrophic gastritis and Helicobacter pylori infection among Japanese Americans in Seattle. *Am J Epidemiol* 2000;151:820–30.
- Selinger CP, Nelson-Piercy C, Fraser A, *et al*. Ibd in pregnancy: recent advances, practical management. *Frontline Gastroenterol* 2021;12:214–24.
- van der Woude CJ, Ardizzone S, Bengtson MB, *et al*. The second European evidenced-based consensus on reproduction and pregnancy in inflammatory bowel disease. *J Crohns Colitis* 2015;9:107–24.
- Chua N, Parker E, Giles I, *et al*. A systematic review of live vaccine outcomes in infants exposed to biologic disease modifying anti-rheumatic medications (DMARDs) in-utero. *Rheumatology* 2022. doi:10.1093/rheumatology/keac141. [Epub ahead of print: 08 Mar 2022].