

# Opioid and benzodiazepine prescription among patients with cirrhosis compared to other forms of chronic disease

Monica A Konerman,<sup>1</sup> Mary Rogers,<sup>2</sup> Brooke Kenney,<sup>2</sup> Amit G Singal,<sup>3</sup> Elliot Tapper,<sup>1</sup> Pratima Sharma,<sup>4</sup> Sameer Saini,<sup>2</sup> Brahmajee Nallamothu,<sup>5</sup> Akbar Waljee<sup>2</sup>

**To cite:** Konerman MA, Rogers M, Kenney B, *et al*. Opioid and benzodiazepine prescription among patients with cirrhosis compared to other forms of chronic disease. *BMJ Open Gastro* 2019;**6**:e000271. doi:10.1136/bmjgast-2018-000271

Received 30 December 2018  
Revised 21 February 2019  
Accepted 23 February 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

<sup>1</sup>Department of Internal Medicine, University of Michigan Hospital, Ann Arbor, Michigan, USA

<sup>2</sup>University of Michigan, Ann Arbor, Michigan, USA

<sup>3</sup>Internal Medicine, University of Texas Southwestern, Dallas, Texas, USA

<sup>4</sup>Division of Gastroenterology, University of Michigan, Ann Arbor, Michigan, USA

<sup>5</sup>VA Center for Clinical Management Research, VA Ann Arbor Health Care System, Ann Arbor, Michigan, USA

## Correspondence to

Dr Monica A Konerman;  
konerman@med.umich.edu

## ABSTRACT

**Objective** Data on patterns and correlates of opioid and benzodiazepines prescriptions among patients with chronic conditions are limited. Given a diminished capacity for hepatic clearance, patients with cirrhosis represent a high risk group for use. The aim of this study was to characterise the patterns and correlates of prescription opioid, benzodiazepine and dual drug prescriptions among individuals with common chronic diseases.

**Design** Analysis of Truven MarketScan database to evaluate individuals with drug coverage with cirrhosis (n=169,181), chronic hepatitis C without cirrhosis (n=210 191), congestive heart failure (n=766 840) or chronic obstructive pulmonary disease (n=1 438 798). Pharmacy files were examined for outpatient prescriptions.

**Results** Patients with cirrhosis had a significantly higher prevalence of opioid prescriptions (37.1 per 100 person-years vs 24.3–26.0, p<0.001) and benzodiazepine prescriptions (21.3 per 100 person-years vs 12.1–12.9, p<0.001). High dose opioid prescription (≥90 daily oral morphine equivalents) (29.1% vs 14.4%, p<0.001) and dual opioid and benzodiazepine prescription (17.5% vs 9.6%–10.5%, p<0.001) were also significantly more prevalent in cirrhosis. High dose opioid prescription was greater in men, individuals ages 40–59, in the Western USA, and among those with a mental health or substance abuse condition. Dual opioid and benzodiazepine prescription were highest among those with alcoholic cirrhosis and middle aged-adults.

**Conclusion** Persons with cirrhosis have significantly higher rates of prescription opioid and benzodiazepine prescription compared to others with chronic diseases despite their high risk for adverse drug reactions. Demographics and mental health or substance abuse history can help identify high risk groups to target interventions.

## INTRODUCTION

Approximately 70.6 opioid prescriptions are filled per 100 persons in the USA each year.<sup>1</sup> The overall rates of opioid prescriptions quadrupled between 1999 and 2015 with annual

## Summary box

### What is already known about this subject?

- ▶ Prescription opioid and benzodiazepine use is highly prevalent and associated with increased morbidity and mortality in the general population.
- ▶ Impact of these medications among patients with chronic liver disease (CLD) may differ due to their hepatic based metabolism.
- ▶ There are very limited data on patterns and correlates of use of prescription opioids and benzodiazepines in chronic conditions including CLD.

### What are the new findings?

- ▶ In this large, nationwide database of insured US patients, we demonstrate that prescription of opioids, benzodiazepines and dual prescription are significantly higher in patients with cirrhosis than in individuals with other common chronic conditions (congestive heart failure and chronic obstructive pulmonary disease).
- ▶ High dose opioid prescription is also most common in cirrhosis and aetiology of liver disease also appears to drive patterns of prescription.

### How might it impact on clinical practice in the foreseeable future?

- ▶ Patients with CLD, particularly those with alcoholic liver disease, are high utilisers of prescription opioids and benzodiazepines.
- ▶ This prescription, particularly at the high doses used in this population, has been associated with higher rates of adverse clinical outcomes and thus targeted interventions to curb use in this high risk population should be implemented.

increases in associated adverse outcomes such as unintentional overdoses.<sup>2</sup> While patterns of opioid use around acute conditions and surgical procedures have been extensively studied, little is known about opioid use among patients with chronic disease. In addition, many patients with chronic opioid



use are also prescribed benzodiazepines, which when prescribed in combination with opioids, can increase the propensity for adverse clinical outcomes.<sup>3,4</sup> Available data have suggested increases in rates of combined opioid and benzodiazepine use over the past few years at the population level.<sup>3</sup> To date, however, there has been little to no characterisation of the prevalence of dual opioid and benzodiazepine use in chronic disease. A patient population at particular risk for harms associated with opioid and benzodiazepine use includes individuals with cirrhosis due to the hepatic-based clearance of both medications. As such, characterisation of rates and correlates of prescription opioid and benzodiazepine prescriptions in this cohort is of particular clinical importance.<sup>5</sup>

Multiple studies have identified liver disease as a primary risk factor for opioid-related overdose, respiratory and central nervous system depression.<sup>6,7</sup> Opioid use among patients with cirrhosis has been associated with higher rates of hospital readmissions, concern for worsening hepatic encephalopathy and higher overall mortality.<sup>8</sup> Despite the potentially higher risk for adverse side effects, in clinical practice patients with cirrhosis may be frequently prescribed these medications. Given the patterns of associated sociodemographic characteristics and comorbid conditions among many individuals with cirrhosis, these factors may also contribute to differential prevalence and patterns of use in this patient population compared with other relevant chronic conditions and thus warrant further investigation.

The true scope of this problem remains unknown however, as the precise prevalence and patterns of opioid and benzodiazepine use in patients with cirrhosis, particularly in comparison to rates seen in other common chronic diseases, has not been clearly characterised to date. Current estimates are highly variable and predominantly generated from small, single-centre retrospective studies. The prevalence rates for prescription opioid use in cirrhosis range from 25% to 77% and encompass heterogeneous patient subgroups including analyses restricted to hospitalised patients.<sup>8-13</sup> Moreover, there has not been a clear evaluation of opioid and benzodiazepine use across different forms of chronic liver disease. As such, the aim of this study is to characterise opioid and benzodiazepine prescription among patients with cirrhosis and compare this with rates and patterns of prescription in other chronic diseases. In addition, we aimed to identify correlates of opioid and benzodiazepine prescription in patients with chronic disease.

## PATIENTS AND METHODS

### Study population

A nationwide longitudinal study was conducted using the 2009–2015 Truven Health MarketScan Commercial Claims and Encounters Database and the Medicare Supplemental Database. These data included patients in all 50 states who received health insurance coverage from a range of employer-based health plans. Deidentified

outpatient pharmacy insurance claims were linked to ascertain medical encounters and medication prescription among insured employees and their dependents. For this study, we included patients with diagnosis codes indicating cirrhosis from any aetiology, chronic hepatitis C (HCV) without cirrhosis, congestive heart failure (CHF) and chronic obstructive pulmonary disease (COPD). Codes used for case identification are listed in the (online appendix 1). For this study, single ICD-9 codes were used to identify patients with cirrhosis based on validated algorithms using this approach that have been shown to have positive predictive values of  $\geq 80\%$ .<sup>14,15</sup> Disease groups were mutually exclusive; there were no overlapping diagnoses across these four main categories of diseases in order to optimise comparison of rates of prescription. All patients included had continuous drug coverage over their inclusion period in the study. This study was determined to be exempt for requiring institutional review board approval and overall complies with ethical aspects of clinical research.

### Study outcomes and other covariates

Pharmacy files were examined for the use of prescription opioids and benzodiazepines. Opioids were identified as prescriptions found in the therapeutic drug class (inclusive list can be found in the Appendix). Daily opioid morphine equivalents (OME) were calculated based on the number of pills prescribed, the strength of the prescription and the morphine-equivalent conversion factor. OME has been recommended by the CDC to assess total opioid burden. High dose prescriptions were of interest and therefore, daily OMEs of 90 or greater were determined for each patient based on CDC categorisation for opioid prescription dosing.<sup>16</sup>

Demographics including age at the time of enrolment, gender and geographic region were included. There were no age restrictions, with age categorised by  $\leq 18$ , 19–39, 40–59 and  $\geq 60$  years. Among individuals with cirrhosis, patients were further subcategorised according to aetiology of liver disease: alcoholic cirrhosis with HCV, alcoholic cirrhosis without HCV, non-alcoholic cirrhosis with HCV and all other types of cirrhosis. Clinical variables of interest included the Charlson comorbidity index, and presence of a mental health or substance abuse diagnosis occurring in the year before the index disease diagnosis (codes used for case identification outlined in online appendix).

### Statistical analysis

Prevalence rates of opioid and benzodiazepine drug prescription (per 100 person-years) were calculated with Poisson exact CIs. The number of people prescribed opioids and benzodiazepines constituted the numerator with person-years of observation as the denominator (ie, enrolment period in healthcare coverage) for each of the disease categories. Median number of high-dose prescriptions ( $\geq 90$  daily OME) were determined, with IQR. Percentages of patients using high-dose prescriptions

by disease category were evaluated by Pearson  $\chi^2$  tests. Correlates of the numbers of high dose ( $\geq 90$  daily OME) opioid prescriptions were assessed using negative binomial regression, offset by the natural log of the years under observation. For correlates of opioid and benzodiazepine prescription, the following variables were assessed: age, gender, region, history of substance abuse or mental health conditions, Charlson comorbidity index and aetiology of cirrhosis (among those with cirrhosis). Alpha was set at 0.05, 2-tailed. Analyses were conducted in Stata/MP 15.1 (College Station, Texas, USA).

## RESULTS

### Study population

In this cohort, there were 169 181 patients with cirrhosis, 210 191 patients with HCV without cirrhosis, 766 840 with CHF and 1 438 798 with COPD. The median age of the overall cohort was 59 years (IQR: 50–70) with 50.7% being females. Both cirrhosis and HCV without cirrhosis

were more prevalent in patients 40–59 years of age (table 1) whereas CHF and COPD occurred more often in older patients. Males were more likely to have HCV without cirrhosis than females, while COPD was more common in females.

### Opioid prescription

The prevalence of opioid prescription among patients with cirrhosis was 37.1 per 100 person-years and was significantly greater than in HCV without cirrhosis, CHF and COPD (table 2). The prevalence was 43% greater than in CHF [rate ratio (RR)=1.43, 95% CI 1.42 to 1.44] and 53% greater than in COPD (RR=1.53, 95% CI 1.52 to 1.54). Median daily OMEs among those with cirrhosis was 40 (IQR: 24–90), and for HCV without cirrhosis was 40 (IQR: 25–90). Among patients with cirrhosis, 29.1% (n=31 861) had high dose opioid prescriptions ( $\geq 90$  daily OME) during the study period compared with 14.4% (n=30 307) of patients with HCV without cirrhosis ( $p < 0.001$ ). Results

**Table 1** Demographic characteristics of the patients

Characteristic		Disease			Total
		Cirrhosis	CHF	COPD	
Age (years)					
≤18	n	2922	10 784	53 409	70 495
	%	1.73	1.41	3.71	2.83
19–39	n	17 437	42 473	89 022	179 214
	%	10.31	5.54	6.19	7.18
40–59	n	101 596	235 725	588 456	1 006 691
	%	60.05	30.74	40.90	40.35
≥60	n	47 226	477 858	707 911	1 238 761
	%	27.91	62.32	49.20	49.65
Sex					
Male	n	94 546	395 130	663 076	1 218 848
	%	55.88	51.53	46.09	48.85
Female	n	74 635	371 710	775 722	1 276 313
	%	44.12	48.47	53.91	51.15
Geographic region					
Northeast	n	32 302	162 386	298 722	518 891
	%	19.09	21.18	20.76	20.80
North Central	n	35 596	208 474	401 632	661 723
	%	21.04	27.19	27.91	26.52
South	n	65 703	269 322	521 118	905 957
	%	38.84	35.12	36.22	36.31
West	n	31 235	113 227	186 374	356 996
	%	18.46	14.77	12.95	14.31
Unknown	n	4345	13 431	30 952	51 594
	%	2.57	1.75	2.15	2.07
Total	n	169 181	766 840	1 438 798	2 495 161
	%	100	100	100	100

CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease.

**Table 2** Prevalence of opioid and benzodiazepine prescription by chronic disease

Category	Cirrhosis	Hepatitis C without cirrhosis	CHF	COPD	P value
<b>All</b>					
Number of patients	169 181	210 191	766 840	1 438 798	
Person-years	294 692	456 363	1 811 106	3 716 909	
<b>Opioids</b>					
Number of patients using opioids 2009–2015	109 444	113 486	470 956	902 192	
Prevalence of opioid prescription per 100 person-years (95% CI)	37.1 (37.0 to 37.3)	24.9 (24.7 to 25.0)	26.0 (25.9 to 26.1)	24.3 (24.2 to 24.3)	<0.001
<b>Benzodiazepine</b>					
Number of patients using benzodiazepine 2009–2015	62 703	56 036	219 258	478 494	
Prevalence of benzodiazepine per 100 person-years (95% CI)	21.3 (21.1 to 21.4)	12.3 (12.2 to 12.4)	12.1 (12.1 to 12.2)	12.9 (12.8 to 12.9)	<0.001
<b>Both opioids and benzodiazepines</b>					
Number of patients using both opioids and benzodiazepine 2009–2015	51 475	43 948	175 153	390 820	
Prevalence of both opioid and benzodiazepine per 100 person-years (95% CI)	17.5 (17.3 to 17.6)	9.6 (9.5 to 9.7)	9.7 (9.6 to 9.7)	10.5 (10.5 to 10.5)	<0.001

CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease.

were similar when analysis was restricted to individuals age 65 or younger in order to address the possibility of difficulty capturing opioid prescriptions in this dataset in patients with dual Medicare eligibility (online supplementary file 1). In terms of type of opioid prescribed, only a very small proportion consisted of buprenorphine (0.08%, 0.66%, 0.03%–0.05%) or methadone (0.49%, 0.81%, 0.25% and 0.33%) among patients with cirrhosis, HCV, CHF and COPD, respectively.

### Benzodiazepine prescription

The prevalence of benzodiazepine prescription among patients with cirrhosis was 21.3 per 100 person-years (table 2). This was significantly greater than benzodiazepine prescription in patients with HCV without cirrhosis (RR=1.73, 95% CI 1.71 to 1.75), CHF (RR=1.76; 95% CI 1.74 to 1.77) or COPD (RR=1.65; 95% CI 1.64 to 1.67).

### Dual opioid and benzodiazepine prescription


The prevalence of dual benzodiazepine and opioid prescription was also greater in patients with cirrhosis compared with the other patient groups ( $p<0.001$ ). Among those with cirrhosis, dual prescription was 17.5% compared with 9.6%, 9.7% and 10.5% among HCV without cirrhosis, CHF and COPD, respectively (table 2). This represents an 81% greater rate than in HCV without cirrhosis (RR=1.81; 95% CI 1.79 to 1.84) and in CHF (RR=1.81; 95% CI 1.79 to 1.82) and a 66% greater prescription than in COPD (RR=1.66; 95% CI 1.65 to 1.68).

The proportion of patients receiving both benzodiazepines and opioids over the study period 2009–2015 is illustrated by age categories in figure 1. Combined drug prescription was greater in ages 30–59 for those

with cirrhosis. Dual drug prescription was quite high in those with HCV without cirrhosis for patients who were 10–19 years of age. For patients with CHF, combined drug prescription was somewhat uniform across ages 30–79. For patients with COPD, combined drug prescription peaked in ages 30–59 years and disproportionately affected older adults.

### DRUG PRESCRIPTION BY TYPE OF CIRRHOSIS

Opioid prescription was rather uniform across all subtypes of cirrhosis, from prevalence rates of 35.5 to 38.0 person-years (table 3). Benzodiazepine prescription was greatest in those alcohol contributing to their liver disease compared with those without alcohol abuse.



**Figure 1** Proportion of patients using both opioids and benzodiazepines, years 2009–2015, by age and type of disease. CHF, congestive heart failure; COPD, chronic obstructive pulmonary disease; HCV, hepatitis C virus.

**Table 3** Prevalence of opioid and benzodiazepine prescription by type of cirrhosis

	Type of cirrhosis			
	Alcoholic cirrhosis with HCV	Alcoholic cirrhosis	HCV cirrhosis	Other cirrhosis
Number of patients	12 296	39 596	31 762	85 527
Person-years	24 603	66 718	59 235	144 135
Prevalence of opioid prescriptions per-100 person-years	37.0 (36.4 to 37.6)	36.8 (36.4 to 37.1)	35.5 (35.1 to 35.9)	38.0 (37.8 to 38.3)
Prevalence of benzodiazepine prescriptions per-100 person-years	22.5 (22.0 to 23.0)	25.6 (25.3 to 26.0)	19.6 (19.2 to 19.9)	19.8 (19.6 to 20.0)
Prevalence of both opioid and benzodiazepine prescriptions per-100 person-years	19.6 (19.1 to 20.1)	19.7 (19.4 to 20.0)	16.3 (16.0 to 16.6)	16.6 (16.4 to 16.7)

HCV, hepatitis C virus.

Similarly, rates of dual opioid and benzodiazepine prescription were highest among those with alcohol as a contributing cause of their cirrhosis.

### CORRELATES OF HIGH DOSE PRESCRIPTION OPIOID PRESCRIPTION

Correlates of opioid prescription among those with liver disease are shown in [table 4](#). High dose opioid prescription was more common in adults ages 40–59 years (for both cirrhosis and HCV without cirrhosis) than at other ages. High dose opioid prescription was 26% greater in men than women for patients with cirrhosis, and 30% greater for those with hepatitis C without cirrhosis. High dose opioid prescription was greater in the western USA for both cirrhosis and hepatitis C without cirrhosis. Having a diagnosis of substance abuse or mental health condition was associated with a 62% greater prevalence of high dose opioid prescription in those with hepatitis C without cirrhosis. For those with cirrhosis, it was associated with a 29% greater prevalence. Individuals with two or more comorbidities were more likely to prescription high dose opioids than those with 0–1 comorbidities. For those with cirrhosis, the subtypes most associated with high dose opioid prescription were alcoholic cirrhosis with HCV and non-alcoholic cirrhosis with HCV.

### DISCUSSION

In a large, nationally representative database, we demonstrated that despite a propensity for adverse drug-related side effects, individuals with cirrhosis had the highest rates of opioid, benzodiazepine and dual opioid and benzodiazepine prescription. Individuals with cirrhosis are being frequently prescribed opioids, and a significant proportion of individuals with cirrhosis are being prescribed opioids at very high doses ( $\geq 90$  OMEs). Among individuals with cirrhosis, those with alcoholic liver disease appeared to have the highest rates of benzodiazepine and dual opioid/benzodiazepine prescription, and those with HCV were most likely to have high dose opioid prescription. Overall, patterns of opioid and benzodiazepine prescription appeared to be correlated

with demographic factors and comorbid conditions, with variable rates based on age, sex, geographic region and the presences of mental health conditions and other substance abuse history.

### Highest rates of opioid, benzodiazepine and dual prescription in cirrhosis

In this study, we have added to the existing literature by confirming in a large, nationally representative database the very high rates of prescription opioid and benzodiazepine use in cirrhosis, occurring in one-third and one-fifth of patients, respectively. Importantly, in this study, we have been able to place these rates and patterns of prescription in context relative other comparator chronic diseases that have similarly high overall health-care utilisation.<sup>17</sup> The reasons for this discrepant patterns of prescription in cirrhosis remain largely unknown and represent an important area for future investigation. Chronic pain is a common symptom across these disease states and thus variable rates of prescription would not be anticipated to reflect the burden of symptomatology.<sup>18–20</sup>

In clinical practice, there remains prevalent misconceptions regarding safety of acetaminophen among individuals with chronic liver disease, and this may in part drive higher use of alternative forms of analgesics, but this has not been definitively shown in prior studies.<sup>21 22</sup> Additionally, this factor alone is unlikely to explain this degree of discrepancy in rates of prescriptions and thus dedicated evaluation of patient and provider perceptions and other factors influencing pain management in cirrhosis is needed. Relatedly, another important finding of note was the rates of high dose ( $\geq 90$  OMEs) opioid prescription among patients with cirrhosis, which occurred in 29% of individuals with cirrhosis. Prevalence of high dose opioid prescription was higher in this cohort compared with a large recent VA cohort where  $>100$  OME occurred in 21% of patients, though the overall prevalence of opioid prescription was much higher in that VA population (77%).<sup>13</sup> Given that it is well established that risk of adverse drug effects parallels OMEs, further understanding of these prescribing patterns is urgently warranted.<sup>23</sup>

**Table 4** Correlates of the number of high dose (>90 daily OME) opioid prescriptions

	Cirrhosis				Hepatitis C without cirrhosis			
	Mean # high dose prescriptions	RR	95% CI	P value	Mean # high dose prescriptions	RR	95% CI	P value
Age (years)								
≤18	4.4	0.46	0.37 to 0.57	<0.001	4.6	0.35	0.29 to 0.41	<0.001
19–39	6.8	1.20	1.12 to 1.28	<0.001	6.6	1.07	1.00 to 1.14	0.051
40–59	8.5	1.32	1.27 to 1.37	<0.001	10.0	1.17	1.11 to 1.23	<0.001
≥60	5.1	1.00	Reference		6.6	1.00	Reference	
Sex								
Male	7.6	1.26	1.21 to 1.30	<0.001	9.3	1.30	1.26 to 1.35	<0.001
Female	7.2	1.00	Reference		8.5	1.00	Reference	
Geographic region								
Northeast	8.2	0.93	0.87 to 0.98	0.008	10.0	0.84	0.79 to 0.89	<0.001
North Central	6.8	0.71	0.67 to 0.75	<0.001	8.3	0.69	0.65 to 0.73	<0.001
South	6.8	0.65	0.62 to 0.69	<0.001	8.1	0.70	0.67 to 0.74	<0.001
West	8.7	1.00	Reference		10.1	1.00	Reference	
Substance abuse/mental health diagnosis								
Yes	9.6	1.29	1.24 to 1.35	<0.001	10.8	1.62	1.55 to 1.69	<0.001
No	6.6	1.00	Reference		8.1	1.00	Reference	
Charlson comorbidity index								
2 or higher	7.3	1.25	1.20 to 1.29	<0.001	9.8	1.13	1.08 to 1.19	<0.001
0–1	7.5	1.00	Reference		8.8	1.00	Reference	
Aetiology of liver disease								
Alcoholic cirrhosis with HCV	9.7	1.25	1.17 to 1.33	<0.001	N/A			
Alcoholic cirrhosis (without HCV)	7.2	0.98	0.93 to 1.03	0.353				
HCV cirrhosis (without alcohol)	9.4	1.19	1.13 to 1.25	<0.001				
Other cirrhosis*	6.3	1.00	Reference					

\*Includes all forms of cirrhosis excluding HCV and alcoholic related cirrhosis.

HCV, hepatitis C virus; N/A, not applicable; OME, opioid morphine equivalents; RR, relative risk.

### Variable patterns of prescription by demographics and comorbidities

Another key finding of our study is the identification of correlates of opioid and benzodiazepine prescription among individuals with chronic disease. Middle-aged individuals had significantly higher rates of high dose opioid prescription as well as dual opioid-benzodiazepine prescription. Men were also noted to have much higher rates of opioid prescription. This variation in terms of age and sex has been inconsistently shown in prior studies.<sup>24 25</sup> Differences in risk profiles according to these demographic features may be explained by subgroups of patients evaluated. Of note, in this study, we also found significant differences in rates of prescription according to geographic region, with highest rates

of opioid prescription among individuals in the western USA. Geographic variations in opioid prescribing in the USA has been noted prior, particularly as it relates to disproportionate prescription in southern and western states, with the reasons for this variation remaining unclear.<sup>1 26</sup>

As expected, among individuals with cirrhosis and HCV without cirrhosis, those with a higher overall disease burden as indicated by a higher Charlson Comorbidity Index had a RR of 1.25 for high dose opioid prescription. Presence of a mental health or other substance abuse condition was particularly high risk (RR 1.29) in terms of likelihood of high dose opioid prescription. This higher rate of opioid prescription among individuals with other mental health conditions has been well described in

the general population, with prior retrospective and cross-sectional studies highlighting the disproportionate prescription of opioids among patients with mood and anxiety disorders.<sup>27 28</sup>

### Patterns of prescription by aetiology of liver disease

A finding of particular interest was the variations in opioid and benzodiazepine prescription by aetiology of cirrhosis. Rates of opioid prescription were generally similar across different aetiologies of cirrhosis, but high dose opioid prescription was more common among patients with HCV as a cause of their cirrhosis. Interestingly, benzodiazepine prescription was markedly different across aetiologies of cirrhosis, with individuals with alcohol contributing to their cirrhosis having the highest rates of benzodiazepine and dual opioid and benzodiazepine prescription (22%–25% vs 19% and 19% vs 16%, respectively). This type of examination of rates of opioid and benzodiazepine prescription by type of liver disease has not been previously described. It has been hypothesised based on observations in clinical practice that individuals with prior alcohol use disorders increase uptake of benzodiazepines, in part due to counselling about need for alcohol cessation.<sup>29</sup> This change in use of substances is particularly relevant among individuals with liver disease whose access to liver transplantation is restricted based on alcohol use and thus may drive increased use of alternative agents with neuropsychiatric effect, though this concept has not been the subject of dedicated investigation.

### Contextual factors and limitations of this study

There are several limitations to note related to this study. The primary limitation stems from the use of a secondary database which relies on diagnosis codes for categorisation of chronic disease. There is inherent potential for misclassification with both undercoding and overcoding. Marketscan has been analysed extensively for the purpose of clinical research and diagnosis codes for these common conditions have been applied in multiple studies for evaluation of outcomes in chronic disease. Another point to highlight includes the lack of evaluation of inpatient or emergency room prescriptions and potential inciting events (such as surgeries or other acute inciters including delirium tremens) and their impact on overall prevalence of use. Similarly, given the nature of this dataset, we were only able to investigate a limited set of variables as correlates of opioid and benzodiazepine prescription.

### Clinical and health policy implications and conclusion

In conclusion, we demonstrated high rates of prescription opioid, benzodiazepine and dual opioid and benzodiazepines among individuals with chronic disease, with markedly higher rates of prescription among individuals with cirrhosis. Rates of prescriptions are high in cirrhosis, and dosages among patients with cirrhosis are also markedly higher, further compounding potential for adverse

clinical outcomes in this high risk patient population. Rates of prescription varied by age, sex geographic region and aetiology of liver disease highlighting important variations in prescribing patterns that may be amenable to targeted interventions.

**Contributors** MAK: study design and drafting of manuscript. MR: statistical analysis. BK: statistical analysis and revision of manuscript. AGS: study design, revision of manuscript. ET: revision of manuscript. PS: revision of manuscript. SS: revision of manuscript. BN: study design and revision of manuscript. AW: study design and revision of manuscript.

**Funding** AW's research is funded by a VA HSR&D CDA-2 Career Development Award (1K2HX000775). AW and BN also receive support from Michigan Institute for Data Science (MIDAS) at the University of Michigan. AGS's research is funded by R01MD12565 and R01 CA12008.

**Disclaimer** These funding sources did not have any role in the design, conduct, collection, management, analysis, interpretation, preparation, review or approval of the manuscript. This content is solely the responsibility of the authors and does not necessarily represent the official views of the healthcare centres, the NIH or the VA.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data may be obtained from a third party (Truven Marketscan) and are not publicly available. Data are deidentified.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

### REFERENCES

- Guy GP, Zhang K, Bohm MK, *et al*. Vital signs: changes in opioid prescribing in the United States, 2006–2015. *MMWR Morb Mortal Wkly Rep* 2017;66:697–704.
- United States Department of Health and Human Services. The US Opioid epidemic, 2018. Available: <https://www.hhs.gov/opioids/about-the-epidemic/index.html>
- Sun EC, Dixit A, Humphreys K, *et al*. Association between concurrent use of prescription opioids and benzodiazepines and overdose: retrospective analysis. *BMJ* 2017;356.
- Park TW, Saitz R, Ganoczy D, *et al*. Benzodiazepine prescribing patterns and deaths from drug overdose among US veterans receiving opioid analgesics: case-cohort study. *BMJ* 2015;350:h2698.
- Tegeder I, Lotsch J, Geisslinger G. Pharmacokinetics of opioids in liver disease. *Clinical Pharmacokinetics* 1999;37:17–40.
- Zedler B, Xie L, Wang L, *et al*. Development of a risk index for serious prescription opioid-induced respiratory depression or overdose in veterans' health Administration patients. *Pain Medicine* 2015;16:1566–79.
- Zedler B, Xie L, Wang L, *et al*. Risk factors for serious prescription opioid-related toxicity or overdose among Veterans Health Administration patients. *Pain Med* 2014;15:1911–29.
- Acharya C, Betrapally NS, Gillevet PM, *et al*. Chronic opioid use is associated with altered gut microbiota and predicts readmissions in patients with cirrhosis. *Aliment Pharmacol Ther* 2017;45:319–31.
- Rogal SS, Winger D, Bielefeldt K, *et al*. Healthcare utilization in chronic liver disease: the importance of pain and prescription opioid use. *Liver Int* 2013;33:1497–503.
- Rogal SS, Winger D, Bielefeldt K, *et al*. Pain and opioid use in chronic liver disease. *Dig Dis Sci* 2013;58:2976–85.
- Scaglione SJ TR, Mumtaz S, *et al*. Opioid use is associated with all-cause and sepsis-related death among patients with cirrhosis. *Hepatology* 2016;63:1020A.
- Whitehead AJ, Dobscha SK, Morasco BJ, *et al*. Pain, substance use disorders and opioid analgesic prescription patterns in veterans with hepatitis C. *J Pain Symptom Manage* 2008;36:39–45.
- Rogal SS, Beste LA, Youk A, *et al*. Characteristics of Opioid Prescriptions to Veterans With Cirrhosis. *Clin Gastroenterol Hepatol*



2018. doi:10.1016/j.cgh.2018.10.021. [Epub ahead of print: 17 Oct 2018].
14. Mellinger JL, Shedden K, Winder GS, *et al*. The high burden of Alcoholic cirrhosis in privately insured persons in the United States. *Hepatology* 2018;68:872–82.
  15. Nehra MS, Ma Y, Clark C, *et al*. Use of administrative claims data for identifying patients with cirrhosis. *J Clin Gastroenterol* 2013;47:e50–4.
  16. Dowell DHT, Chou R. CDC guideline for prescribing opioids for chronic pain. *MMWR Recomm Rep* 2016;65:1–49.
  17. Asrani SK, Kouznetsova M, Ogola G, *et al*. Increasing health care burden of chronic liver disease compared with other chronic diseases, 2004–2013. *Gastroenterology* 2018;155:719–29.
  18. Loftis JM, Hauser P. Pain and opioid use in chronic liver disease: optimal treatment must address the mental health care needs of the patient. *Digestive Diseases and Sciences* 2013;58:2753–5.
  19. Roberts MH, Mapel DW, Hartry A, *et al*. Chronic pain and pain medication use in chronic obstructive pulmonary disease. A cross-sectional study. *Ann Am Thorac Soc* 2013;10:290–8.
  20. Gan Q, Zhang F-ru, Zhou Q-fen, *et al*. Clinical significance of pain in patients with chronic heart failure. *Chin Med J* 2012;125:3223–7.
  21. Chandok N, Watt KDS. Pain management in the cirrhotic patient: the clinical challenge. *Mayo Clinic Proceedings* 2010;85:451–8.
  22. Imani F, Motavaf M, Safari S, *et al*. The therapeutic use of analgesics in patients with liver cirrhosis: a literature review and evidence-based recommendations. *Hepatitis Monthly* 2014;14:e23539.
  23. Els C, Jackson TD, Kunyk D, *et al*. Adverse events associated with medium- and long-term use of opioids for chronic non-cancer pain: an overview of Cochrane reviews. *The Cochrane database of systematic reviews* 2017;10.
  24. Back SE, Payne RL, Wahlquist AH, *et al*. Comparative profiles of men and women with opioid dependence: results from a national multisite effectiveness trial. *Am J Drug Alcohol Abuse* 2011;37:313–23.
  25. Centers for Disease Control and Prevention. CDC, opioid overdose, data. *Prescribing Data* <https://www.cdc.gov/drugoverdose/data/prescribing.html>
  26. McDonald DC, Carlson K, Izrael D. Geographic variation in opioid prescribing in the U.S. *J Pain* 2012;13:988–96.
  27. Davis MA, Lin LA, Liu H, *et al*. Prescription opioid use among adults with mental health disorders in the United States. *J Am Board Fam Med* 2017;30:407–17.
  28. Seal KH, Shi Y, Cohen G, *et al*. Association of mental health disorders with prescription opioids and high-risk opioid use in US veterans of Iraq and Afghanistan. *JAMA* 2012;307:940–7.
  29. Stein MD, Kanabar M, Anderson BJ, *et al*. Reasons for benzodiazepine use among persons seeking opioid detoxification. *J Subst Abuse Treat* 2016;68:57–61.