Traditional serrated adenoma: an overview of pathology and emphasis on molecular pathogenesis

Aoife J McCarthy, Stefano Serra, Runjan Chetty[✉]

To cite: McCarthy AJ, Serra S, Chetty R. Traditional serrated adenoma: an overview of pathology and emphasis on molecular pathogenesis. *BMJ Open Gastro* 2019;6:e000317. doi:10.1136/bmjgast-2019-000317

Received 3 June 2019

Revised 2 July 2019

Accepted 8 July 2019

ABSTRACT

Objective To provide an overview of the pathology and molecular pathogenesis of traditional serrated adenomas (TSA).

Design Describe the morphology and molecules that play a role in their pathogenesis.

Results These exuberant polypoid lesions are typified by tall cells with deeply eosinophilic cytoplasm, elongated nuclei bearing delicate chromatin, ectopic crypt foci, deep cleaving of the lining mucosa and an overall resemblance to small bowel mucosa.

Broadly, TSAs arise via three mechanisms. They may be *BRAF* mutated and CpG island methylator phenotype (CIMP)-high: right sided, mediated through a microvesicular hyperplastic polyp or a sessile serrated adenoma, may also have *RNF43* mutations and result in microsatellite stable (MSS) colorectal cancers (CRC). The second pathway that is mutually exclusive of the first is mediated through *KRAS* mutation with CIMP-low TSAs. These are left-sided TSAs, are not associated with another serrated polyp and result in MSS CRC. These TSAs also have *RSPO3*, *RNF43* and *p53* mutations together with aberrant nuclear localisation of β -catenin. Third, there is a smaller group of TSAs that are *BRAF* and *KRAS* wild type and arise by as yet unknown molecular events. All TSAs show retention of mismatch repair proteins.

Conclusion These are characteristic unusual polyps with a complex molecular landscape.

INTRODUCTION

Recent investigations into the pathogenesis of colorectal carcinoma (CRC) have identified the serrated pathway of colorectal carcinogenesis, which accounts for 15%–35% of CRCs.^{1–4} There are several potentially different precursor lesions implicated in the serrated pathway that fit into the general category of ‘serrated adenomas’. The term ‘serrated adenoma’ was introduced by Longacre and Fenoglio-Preiser, and later refined initially by Torlakovic *et al*, and then by others, to incorporate a spectrum of lesions, namely hyperplastic polyps (HP), sessile serrated adenomas/polyps (SSA; also called sessile serrated lesions in the UK and

parts of Europe), and traditional serrated adenomas (TSA).^{1 5–9}

When TSAs were first described by Longacre and Fenoglio-Preiser, they described mixed hyperplastic adenomatous polyps/serrated adenomas.^{5 10} The authors appreciated that, when compared with HPs, these polyps had certain differences, namely prominent nucleoli, goblet cell immaturity, and absence of a thickened basement membrane.⁵ As awareness of the characteristic morphology of TSAs grew, they are now considered the most unique and easily identified of the serrated lesions.¹¹

The origin of TSAs is unclear, but some probably arise *de novo*, while many appear to arise in a precursor polyp, typically microvesicular HP or SSA.^{1 12–19}

The purpose of this review is to provide a brief clinicopathological background of TSAs and present the current state of knowledge regarding their molecular pathogenesis.

Clinical and epidemiological features

TSAs account for <1% of all colorectal polyps in most series, and for 1%–7% of all serrated lesions.^{5 20–24} TSAs tend to occur in older patients (usually over 50 years of age) and have no significant gender predilection.^{12 15 17} They are found predominantly in the distal (left) colon, and occur only rarely in the proximal colon.¹ Most are less than 10 mm in size.

Morphology

In short, the morphological criteria for diagnosing a TSA include typical cytology (ie, elongated, narrow pencillate nuclei with delicate dispersed chromatin and cytoplasmic hypereosinophilia), ectopic crypt foci (ECF), and typical slit-like clefted serration.⁶ At least two of these three features are required (with at least one of these features being present in 50% of the polyp) to render a diagnosis of TSA with the slit-like serration being the most consistent histological feature.¹² A

© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Division of Anatomical Pathology, Laboratory Medicine Program, University Health Network and University of Toronto, Toronto, Ontario, Canada

Correspondence to

Dr Runjan Chetty;
runjan.chetty@gmail.com

reproducibility study identified this lesion as the one with the best kappa statistic of the various serrated lesions.²⁵

Architecture

TSAs have an overall protuberant exophytic configuration, with a complex villous growth pattern. Mucosal protrusions, in the form of a 'tennis-racquet-like' enlargement of the tip of a protrusion, have been described as a feature unique to TSA.¹

Some TSAs demonstrate a flat growth pattern ('flat' TSA),¹ with the majority of these polyps being elevated less than twice the height of the normal mucosa and lacking prominent villiform projections.¹² They are typically found in the proximal colon,¹² and Bettington and colleagues demonstrated that flat TSAs can be reliably distinguished from SSAs.¹²

Cytological features

The lesional cells have centrally placed, elongated, penicillate nuclei (typically not hyperchromatic) (figure 1A).³⁵ These lesions are usually not mitotically active, as determined by mitotic count or Ki67 immunoreactivity.^{11 26}

The authors have appreciated that the lesional cells in TSAs more or less resemble the normal mucosa of the duodenum, having a brush-border, indented, flat-topped luminal slit-like serrations, pencillate nuclei and eosinophilic cytoplasm.²⁷

Serrations

Slit-like serrations refer to narrow slits in the deeply eosinophilic epithelium similar to normal small intestinal

mucosa.⁴ It has been suggested that these slit-like serrations (figure 1B) are possibly the most specific feature when faced with the challenge of differentiating a TSA from a morphologically similar tubulovillous adenoma with prominent serration.²⁸

Ectopic crypt foci

ECFs refer to the abnormal development of crypts with loss of orientation towards the muscularis mucosae.²⁹ These ectopic crypts are not present in all areas of all TSAs but should be present at least focally, and, in fact, it is thought that they are probably the best defining feature of TSAs (figure 1C), although most TSAs will also have the atypical eosinophilic cell.^{13 13}

Once ECFs and slit-like serrations become identifiable in a small (<10 mm) polyp, the diagnosis of a TSA should be considered, even if villous change is not apparent.³⁰

Other features

Goblet cells are not infrequent in TSAs and their amount varies greatly from case to case, as well as from area to area in the same polyp. Some TSAs are composed mainly of mucin-filled goblet cells ('mucin-rich TSA'),^{31 32} and these mucin-rich TSAs have been arbitrarily defined by the presence of 50% or more goblet/mucin-rich cells with a goblet cell:eosinophilic, absorptive cell ratio of at least 1:1 (31) (figure 1D).

TSAs have been shown to have a relatively increased number of intraepithelial lymphocytes (figure 1E), as compared with classical adenomas, but much less than that of SSAs harbouring conventional adenomatous dysplasia.²⁷ Furthermore, the mucin-rich TSAs have been shown to have more intraepithelial lymphocytes than classic TSAs.³¹

TSAs that occur in the right colon tend to occur more frequently in females, tend to be more of the mucin-rich variety and have greater intraepithelial lymphocytes than their left-sided counterparts.³¹ Furthermore, right-sided TSAs are usually BRAF mutated and have methylation of CpG islands (CpG island methylator phenotype (CIMP) positive)—see molecular discussion below.

Precursor polyp

Many TSAs contain adjacent areas of HP (characterised by glands with saw-toothed, serrated luminal profiles and straight or V-shaped basal or lower one-third gland profiles) or SSA (glands which are also saw-toothed but with very characteristic basal architecture: instead of being straight the basal aspects of glands are dilated, fan out horizontally and give rise to boot-shaped or L-shaped glandular profiles), thought to be a precursor polyp in this setting^{12 17 27 33} (figure 1F). In a study exploring the association of precursor polyps with TSA, 28 were HPs (36%) and 18 were SSAs (23%).³³ Thus, up to one-third of TSAs may contain histological evidence of an HP or SSA suggesting a relationship among all three types of serrated polyps. Evidence of one of these precursor components is characterised by a discrete area of the

Figure 1 The lesional cells of a traditional serrated adenoma (TSA) have centrally placed, elongated, penicillate nuclei, with dispersed chromatin (A). Narrow slits in the epithelium ('slit-like serrations') similar to normal small intestinal mucosa (abundant eosinophilic/oncocyctic cytoplasm) are possibly the most specific feature of a TSA (B). Ectopic crypt foci (ECF) refer to the abnormal development of crypts with loss of orientation towards the muscularis mucosae (C). 'Mucin-rich TSAs' have 50% or more goblet/mucin-rich cells, with a goblet cell:eosinophilic, absorptive cell ratio of at least 1:1 (D). TSAs have a relatively increased number of intraepithelial lymphocytes (arrows) (E). Many TSAs (black circle) contain adjacent areas of hyperplastic polyp or sessile serrated adenoma (SSA) (red circle), with these latter components being thought to represent a precursor polyp in this setting (F).

lesion with clear morphological distinction from the TSA component, either at the edge of, or underlying, the TSA.¹⁵

Dysplasia

Although some authors interpret the prototypical lesion cell of TSA as being inherently dysplastic, others maintain that this cell type itself is generally non-proliferative and therefore may, in fact, represent a metaplastic or a senescent cell.^{1 3 4 10–12 18 19 26 34–42}

The presence of an abrupt transition from typical TSA to discrete areas of cytological atypia or dysplasia of a more conventional type can be seen with varying frequency. The conventional dysplasia is often found towards the base of the TSA,^{1 3 15 43 44} and it has been proposed that these lesions be referred to as 'TSAs with conventional cytological dysplasia',³ or 'advanced TSAs'.¹² Bettington and colleagues showed that advanced TSAs tend to be larger than ordinary TSAs.¹²

Conventional dysplasia

Cytological features of conventional adenomatous dysplasia include increased nuclear size, nuclear crowding, hyperchromasia, complete loss of polarity, and pseudostratification with nuclei extending into the upper half of the neoplastic cell, in addition to frequent and atypical mitoses.^{35 43 44} Architectural features of conventional adenomatous dysplasia include crowding of glands, cribriform glands, and intraluminal necrosis.³⁵

Serrated dysplasia

Serrated dysplasia, as defined by the WHO, is characterised by cuboidal cells with eosinophilic cytoplasm, vesicular nuclei and prominent nucleoli.³⁵

For now, assigning a grade of dysplasia or dividing dysplasia into serrated versus conventional types has no clinical utility and the practising gastroenterologist should not treat a TSA with low-grade dysplasia any differently to a TSA without overt dysplasia and one-off surveillance colonoscopy at 3 years should be performed.^{12 45}

Biomarker expression

TSAs do not express a wide range of routine biomarkers and there are no specific markers available yet. With regard to some of the newer biomarkers such as annexin A10, Hairy and Enhancer of Split 1 (Hes-1) and SPARC-related Modular Calcium-binding 1 protein (SMOC1) more studies are required to confirm their applicability in TSAs.

Ki-67 and CK20

In TSAs, Ki-67 expression is limited to the ECF and basal aspects of the crypts. The opposite staining pattern is found with CK20 where expression is almost exclusively limited to the superficial lining of the surface of the polyp, without extension into the budding crypts.^{1 12}

Annexin A10 and Hes-1

Recent studies have shown that expression of annexin A10 is a reliable marker of SSAs and the serrated pathway of CRC,^{16 17 46–51} and loss of Hes-1 expression can reliably differentiate SSAs from HPs.⁵² Nourbakhsh and Minoo sought to interrogate the concept that at least some TSAs may arise in association with precursor HP or SSA lesions, particularly those that develop in the right colon, by applying these two stains to a series of polyps from the right side of the colon with mixed features of TSA and SSA (as defined by Rex *et al*).^{7 16} The TSA components of these 'hybrid or mixed polyps' showed a staining pattern similar to that of SSAs (annexin A10 overexpression and Hes-1 loss), which the authors suggest supports the theory that SSAs are precursor lesions for at least some TSAs.¹⁶

SPARC-related Modular Calcium-binding 1 protein

Aoki *et al* showed that SMOC1 is expressed immunohistochemically (cytoplasmic staining) in normal colonic epithelium and in SSAs, but its expression is decreased in TSAs.⁵³ Thus, they suggested that immunohistochemical staining of SMOC1 is highly discriminative between TSAs and SSAs, and could be used as an ancillary tool in challenging polyps. They also demonstrated that *SMOC1* is frequently methylated in TSAs and rarely methylated in SSAs.⁵³

p53/p16/β-catenin

Bettington and colleagues showed that advanced TSAs (defined as those with overt dysplasia or carcinoma) displayed strong p53 staining in just over half of cases,¹² which is an adequate surrogate for *TP53* gene mutation.

They demonstrated nuclear β-catenin staining in 40% of cases, indicative of Wnt pathway activation as an important step in malignant progression.¹²

The authors also showed loss of p16 staining in the advanced areas of just over half of *BRAF*-mutant TSAs, and of one-tenth of *KRAS*-mutant TSAs or *BRAF*/*KRAS*-wild-type TSAs.¹² This loss is attributed to methylation-induced silencing of the *CDKN2A* gene which appears to be an important step in the development of adenocarcinoma in these polyps.¹²

Mismatch repair proteins

Mismatch repair (MMR) enzyme function is retained in effectively all TSAs, even when they develop high-grade dysplasia or invasive adenocarcinoma, implying a microsatellite stable (MSS) phenotype.^{12 44}

MMR deficiency has, however, been demonstrated in TSAs from proven *MMR* gene mutation carriers, with loss of immunohistochemical expression in keeping with the underlying germline mutation.⁵⁴ High-level microsatellite instability (MSI-H) results were shown to be concordant with MMR protein loss results in these polyps.⁵⁴ None of the cases tested (all of which were *MLH1*-deficient polyps) demonstrated *MLH1* methylation or somatic *BRAF*c.1799T>A (p.V600E) mutation.⁵⁴

Very rarely, sporadic TSAs do display *MLH1* hypermethylation.^{12 27} Bettington *et al* showed in their series of 200 TSAs that *MLH1* promoter methylation was present in 7% of the *BRAF*-mutant TSAs (eight ordinary/classical and one advanced), but in none of the *KRAS*-mutant, or *BRAF/KRAS*-wild-type TSAs.¹² Only the advanced TSA with *MLH1* methylation showed concordant loss of *MLH1* expression by immunohistochemistry.¹²

Molecular changes

As previously stated, it is now known that 15%–35% of CRCs arise as a consequence of the serrated neoplasia pathway.^{2–4} Mitogen-activated protein kinase (MAPK) pathway activation, a critical early event as a consequence of either activating *BRAF* or *KRAS* mutation,⁵⁵ and the CIMP, causing methylation of CpG islands in the promoter regions of several genes resulting in gene silencing, are well-established molecular drivers of this pathway.^{55–57}

TSAs are thought to arise via three molecular pathways.

The first mechanism is via *BRAF* mutation and CpG island methylation (CIMP) resulting in the CIMP-high (CIMP-H) phenotype. These TSAs tend to be right sided, maybe associated with a precursor microvesicular HP, or an SSA. Furthermore, they may also have *RNF43* mutations and result in MSS colorectal cancers (CRC). The second pathway that is mutually exclusive of the first is mediated through *KRAS* mutations and these are CIMP-low (CIMP-L) TSAs. These TSAs are usually located in the left colon, are not associated with another precursor serrated polyp but also result in MSS CRC like the first pathway described above. These TSAs may also show *RSPO3*, *RNF43* and *p53* mutations together with aberrant, nuclear localisation of β -catenin.

The third pathway resulting in a smaller group of TSAs is both *BRAF* and *KRAS* wild type and arise by, as yet, unknown molecular events.

Although TSAs are genetically a heterogeneous group they are recognised as a precursor lesion for MSS or low-level MSI CRC and, only rarely lead to sporadic MSI-H colon cancers.⁴¹⁹ This even applies to those TSAs harbouring a *BRAF* mutation that are mostly located in the right colon.^{12 15 18}

MAPK pathway

Studies have reported either *BRAF* or *KRAS* mutations in the vast majority (>80%) of TSAs,²⁷ and *BRAF* (22%–42% of TSAs) and *KRAS* (48%–67% of TSAs) mutations are mutually exclusive.^{14 15 17 27 37 43 44}

KRAS mutations occur predominantly at codon 12 and less frequently at codon 13, and the most common mutations are *G12D*, *G12V* and *G13D* occurring in 0%–28% of TSAs.^{20 57 58} With regard to *BRAF*, the most frequent mutation is V600E which occurs in 60%–76% of TSA.^{20 57–60}

It has been shown that, although *BRAF*-mutated and *KRAS*-mutated TSAs are molecularly different, these lesions show the same morphology.^{27 44 61} However, *BRAF*-mutant TSAs are more often located in the

proximal colon and are thought to have more frequent origin in an associated precursor polyp than *KRAS*-mutant TSAs.¹² On the other hand, *KRAS*-mutant TSAs are almost exclusively located in the distal colon, especially the rectum.¹² Furthermore, most mucin-rich TSAs, which are typically right sided, also harbour a *BRAF* mutation.³² A small proportion of TSAs are *BRAF/KRAS* wild type, and these cases segregate best with the *KRAS*-mutated group.¹²

CpG island methylation

CIMP-H status has been shown to correlate strongly with both proximal colonic location and *BRAF* mutation.¹²

It has been shown that *BRAF*-mutant TSAs are more often CIMP-H than either *KRAS*-mutant, or *BRAF/KRAS*-wild-type TSAs, while the majority of *KRAS*-mutant, or *BRAF/KRAS*-wild-type TSAs are either CIMP-L or negative for CIMP.^{12 44} Furthermore, TSAs found in the proximal colon are more likely to be CIMP-H, compared with those in the distal colon.^{12 44} Also, in the distal colon, *BRAF*-mutant TSAs are more likely to be CIMP-H than *KRAS*-mutant TSAs.^{12 44 62}

Wnt pathway

Wnt pathway activation, resulting from *Protein Tyrosine Phosphatase, Receptor-type K-R-Spondin 3* (*PTPRK-RSPO3*) gene fusions or *Ring Finger Protein 43* (*RNF43*) gene mutations, occurs in the majority of TSAs.⁶² *RSPO* fusions and *RNF43* mutations are characteristic of TSA and are rare or absent in other types of colorectal polyps.^{63 64}

RSPO fusions

The *PTPRK-RSPO3* fusions in TSAs are mutually exclusive with other Wnt pathway gene alterations, and are thought to be responsible for Wnt pathway activation,⁶³ potentiating ligand-dependent Wnt signalling.⁶⁵ Sekine and colleagues showed that *RSPO* fusions were rarely observed in TSAs located in the proximal colon.⁶⁶ They showed that TSAs with *RSPO* fusions tended to be larger, and to have ECF and a high-grade dysplastic component; while slit-like serrations were less prominent, and associations with precursor polyps being rare in these *RSPO* fusion-positive TSAs.⁶⁵ It has been shown that TSAs with *RSPO* fusions are less likely to have *BRAF* mutations, and more frequently have *KRAS* mutations.^{63 66}

RNF43 mutations

RNF43 mutations have been detected in over 25% of TSAs.^{63 64}

Most of the *RNF43* mutations that have been detected in the TSAs were homozygous, indicating the biallelic inactivation of these tumour suppressor genes.³⁶

RNF43 mutations have also been detected in a small number of TSA-precursor polyps.^{36 63 64 67} It has been demonstrated that *RNF43* mutations in the precursor polyps are heterozygous, in contrast to the homozygous mutations found in the established TSA components,³⁶ suggesting that biallelic inactivation of *RNF43* occurs during the progression to TSAs in these lesions.³⁶

Germline mutations of *RNF43* have been identified in a minority of patients satisfying the WHO criteria of serrated polyposis syndrome in two studies,^{67–69} leading Yan *et al* to suggest that routine germline testing for *RNF43* mutation should be performed in serrated polyposis syndrome families.⁶⁷

TSAs with a precursor component

Molecular studies previously done in both components (ie, on the TSA component and on the HP/SSA component) of TSAs with a histologically identifiable precursor component (HP/SSA) confirmed identical mutations of *KRAS* or *BRAF* in both components, indicating that MAPK pathway gene mutations are the earliest molecular abnormality occurring in the serrated pathway of tumourigenesis.^{14 15 36}

BRAF mutations were more commonly seen in TSAs with an identifiable precursor component located in the right colon, supporting the hypothesis that HPs/SSAs are indeed precursors for TSAs.¹⁴ Thus, it has been suggested that in a *BRAF*-mutated SSA, methylation of *MLH1* takes the oncogenic pathway to MSI-H CRCs, with the SSA progressing to an intermediate stage of SSA with cytological dysplasia before full-blown malignancy.¹⁶ However, it is believed that if Wnt pathway becomes activated (eg, by *RNF43* mutation) before methylation extends to the *MLH1* locus, then the SSA will transform into a TSA and then the lesion will progress through the MSS pathway and almost never goes back to being MSI-H.^{16 63 64 67 70}

Miscellaneous genes

A variety of other changes at the molecular level have also been identified in a subset of TSAs. While further studies are required to confirm their roles, these include:

p16 hypermethylation

p16 hypermethylation has been shown to be characteristic of TSAs.²⁷

GNAS mutation

GNAS mutation has been identified in a small fraction of TSAs (<10%), mainly with concomitant *BRAF* mutation,^{17 63 71} but *GNAS* mutation status does not correlate with advanced histology (ie, TSAs with high-grade dysplasia and/or invasive carcinoma).^{17 71 72} Liu and colleagues concluded in their study that no histological features separate *GNAS*-mutant TSAs from *GNAS*-wild-type TSAs.⁷¹

Others

Mutations in *APC* or *CTNGB1* are limited to a minor subset of TSAs.^{12 43} In addition, *GREM1* and its protein are thought to play a role in TSAs and are highly expressed especially in the ECFs of TSAs.⁷³

Significance

There is evidence that some of these serrated lesions lead to certain subtypes of CRC, which account for interval carcinomas found during endoscopic surveillance

Figure 2 Schematic representation of the putative molecular pathogenesis of TSA showing three pathways. Purple areas represent dysplastic foci. CIMP-H, CpG island methylator phenotype-high; MMR, mismatch repair; MSS, microsatellite stable; SSA, sessile serrated adenoma; TSA, traditional serrated adenoma.

programmes and which are biologically different from the classical Vogelstein model for CRC.^{27 74} A 3-year colonoscopy surveillance interval is currently recommended by many guidelines for usual TSA.^{75–79}

A carcinoma arising in TSA can be the typical colorectal adenocarcinoma not otherwise specified, but can also be a mucinous or serrated adenocarcinoma.³⁹ They are often CIMP-H, but are essentially never MSI.

CONCLUSION

TSA has a characteristic constellation of morphological features that make microscopic recognition of classic examples relatively simple. Histological variants have been described and maintain the basic histological tenets seen in typical TSA. The relative simplicity of morphology is counterbalanced by the molecular complexity that TSAs manifest (figure 2). As we have highlighted, despite the fundamental dichotomy of *BRAF* versus *KRAS*-mutated TSA, the morphological appearances remain the same. At this juncture, there are no compelling reasons to believe that the different molecular pathways confer varying propensities for the development of carcinoma.

Contributors The three authors (AJMC, SS, RC) are authors based on the following four criteria applying to all three authors: substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work: AJMC, SS, RC; drafting the work or revising it critically for important intellectual content: AJMC, SS, RC; final approval of the version to be published: AJMC, SS, RC; agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: AJMC, SS, RC. All individuals listed as coauthors of the manuscript (AJMC, SS, RC) qualify for every one of the four criteria listed above.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Torlakovic EE, Gomez JD, Driman DK, *et al.* Sessile serrated adenoma (SSA) vs. traditional serrated adenoma (TSA). *Am J Surg Pathol* 2008;32:21–9.
2. Leggett B, Whitehall V. Role of the serrated pathway in colorectal cancer pathogenesis. *Gastroenterology* 2010;138:2088–100.
3. Snover DC. Update on the serrated pathway to colorectal carcinoma. *Hum Pathol* 2011;42:1–10.
4. Bettington M, Walker N, Clouston A, *et al.* The serrated pathway to colorectal carcinoma: current concepts and challenges. *Histopathology* 2013;62:367–86.
5. Longacre TA, Fenoglio-Preiser CM. Mixed hyperplastic adenomatous polyps/serrated adenomas. A distinct form of colorectal neoplasia. *Am J Surg Pathol* 1990;14:524–37.
6. Snover DC, Ahnen DJ, Burt RW, *et al.* Serrated polyps of the colon and rectum and serrated polyposis. In: Bosman FT, Carneiro F, Hruban RH, *et al.* eds. *Who classification of tumours of the digestive system*. IARC press. Lyon, France, 2010: 160–5.
7. Rex DK, Ahnen DJ, Baron JA, *et al.* Serrated lesions of the colorectum: review and recommendations from an expert panel. *American Journal of Gastroenterology* 2012;107:1315–29.
8. Ensari A, Bilezikci B, Carneiro F, *et al.* Serrated polyps of the colon: how reproducible is their classification? *Virchows Arch* 2012;461:495–504.
9. Rau TT, Agaimy A, Gehoff A, *et al.* Defined morphological criteria allow reliable diagnosis of colorectal serrated polyps and predict polyp genetics. *Virchows Arch* 2014;464:663–72.
10. Choi E-YK, Appelman HD. A historical perspective and Exposé on serrated polyps of the colorectum. *Arch Pathol Lab Med* 2016;140:1079–84.
11. Torlakovic E, Skovlund E, Snover DC, *et al.* Morphologic reappraisal of serrated colorectal polyps. *Am J Surg Pathol* 2003;27:65–81.
12. Bettington ML, Walker NI, Rosty C, *et al.* A clinicopathological and molecular analysis of 200 traditional serrated adenomas. *Mod Pathol* 2015;28:414–27.
13. Yantiss RK, Oh KY, Chen Y-T, *et al.* Filiform serrated adenomas: a clinicopathologic and immunophenotypic study of 18 cases. *Am J Surg Pathol* 2007;31:1238–45.
14. Kim K-M, Lee EJ, Kim Y-H, *et al.* Kras mutations in traditional serrated adenomas from Korea herald an aggressive phenotype. *Am J Surg Pathol* 2010;34:1–75.
15. Kim M-J, Lee E-J, Suh J-P, *et al.* Traditional serrated adenoma of the colorectum: clinicopathologic implications and endoscopic findings of the precursor lesions. *Am J Clin Pathol* 2013;140:898–911.
16. Nourbakhsh M, Minoo P, Annexin A10 and HES-1 immunohistochemistry in right-sided traditional serrated adenomas suggests an origin from sessile serrated adenoma. *Appl Immunohistochem Mol Morphol* 2019. doi:10.1097/PAI.0000000000000740. [Epub ahead of print: 16 Jan 2019].
17. Wiland HO, Shadrach B, Allende D, *et al.* Morphologic and molecular characterization of traditional serrated adenomas of the distal colon and rectum. *Am J Surg Pathol* 2014;38:1–7.
18. Chetty R. Traditional serrated adenoma (TSA): morphological questions, queries and quandaries. *J Clin Pathol* 2016;69:6–11.
19. Tsai J-H, Cheng C-H, Chen C-C, *et al.* Traditional serrated adenoma with BRAF mutation is associated with synchronous/metachronous BRAF -mutated serrated lesions. *Histopathology* 2016;68:810–8.
20. Spring KJ, Zhao ZZ, Karamatic R, *et al.* High prevalence of sessile serrated adenomas with BRAF mutations: a prospective study of patients undergoing colonoscopy. *Gastroenterology* 2006;131:1400–7.
21. Carr NJ, Mahajan H, Tan KL, *et al.* Serrated and non-serrated polyps of the colorectum: their prevalence in an unselected case series and correlation of BRAF mutation analysis with the diagnosis of sessile serrated adenoma. *J Clin Pathol* 2009;62:516–8.
22. Bettington M, Walker N, Rosty C, *et al.* Critical appraisal of the diagnosis of the sessile serrated adenoma. *Am J Surg Pathol* 2014;38:158–66.
23. Higuchi T, Sugihara K, Jass JR. Demographic and pathological characteristics of serrated polyps of colorectum. *Histopathology* 2005;47:32–40.
24. Lash RH, Genta RM, Schuler CM. Sessile serrated adenomas: prevalence of dysplasia and carcinoma in 2139 patients. *J Clin Pathol* 2010;63:681–6.
25. Farris AB, Misdraji J, Srivastava A, *et al.* Sessile serrated adenoma: challenging discrimination from other serrated colonic polyps. *Am J Surg Pathol* 2008;32:30–5.
26. Snover DC, Jass JR, Fenoglio-Preiser C, *et al.* Serrated polyps of the large intestine: a morphologic and molecular review of an evolving concept. *Am J Clin Pathol* 2005;124:380–91.
27. Rau TT, Atreya R, Aust D, *et al.* Inflammatory response in serrated precursor lesions of the colon classified according to who entities, clinical parameters and phenotype-genotype correlation. *J Pathol* 2016;2:113–24.
28. Hafezi-Bakhtiari S, Wang LM, Colling R, *et al.* Histological overlap between colorectal villous/tubulovillous and traditional serrated adenomas. *Histopathology* 2015;66:308–13.
29. Haramis A-P, *et al.* De novo crypt formation and juvenile polyposis on BMP inhibition in mouse intestine. *Science* 2004;303:1684–6.
30. Bettington M, Rosty C, Whitehall V, *et al.* A morphological and molecular study of proposed early forms of traditional serrated adenoma. *Histopathology* 2018;73:1023–9.
31. N Kalimuthu S, Serra S, Hafezi-Bakhtiari S, *et al.* Mucin-rich variant of traditional serrated adenoma: a distinct morphological variant. *Histopathology* 2017;71:208–16.
32. Hiromoto T, Murakami T, Akazawa Y, *et al.* Immunohistochemical and genetic characteristics of a colorectal mucin-rich variant of traditional serrated adenoma. *Histopathology* 2018;73:444–53.
33. Chetty R, Hafezi-Bakhtiari S, Serra S, *et al.* Traditional serrated adenomas (TSAs) admixed with other serrated (so-called precursor) polyps and conventional adenomas: a frequent occurrence. *J Clin Pathol* 2015;68:270–3.
34. Snover DC. Serrated polyps of the large intestine. *Semin Diagn Pathol* 2005;22:301–8.
35. Bosman FT, Carneiro F, Hruban RH, *et al.* International Agency for Research on Cancer. In: *Who classification of tumours of the digestive system*. 4th edn. Lyon: International Agency for Research on Cancer, 2010.
36. Hashimoto T, Ogawa R, Yoshida H, *et al.* Acquisition of Wnt pathway gene alterations coincides with the transition from precursor polyps to traditional serrated adenomas. *Am J Surg Pathol* 2019;43:132–9.
37. Bettington ML, Chetty R. Traditional serrated adenoma: an update. *Hum Pathol* 2015;46:933–8.
38. Leedham SJ, Chetty R. Wnt disruption in colorectal polyps – the traditional serrated adenoma enters the fray. *J Pathol* 2016;239:387–90.
39. O'Brien MJ, Zhao Q, Yang S. Colorectal serrated pathway cancers and precursors. *Histopathology* 2015;66:49–65.
40. Yang H-M, Mitchell JM, Sepulveda JL, *et al.* Molecular and histologic considerations in the assessment of serrated polyps. *Arch Pathol Lab Med* 2015;139:730–41.
41. Bateman AC, Shepherd NA. UK guidance for the pathological reporting of serrated lesions of the colorectum. *J Clin Pathol* 2015;68:585–91.
42. McCarthy AJ, O'Reilly SM, Shanley J, *et al.* Colorectal serrated neoplasia: an institutional 12-year review highlights the impact of a screening programme. *Gastroenterol Res Pract* 2019;2019:1–9.
43. Fu B, Yachida S, Morgan R, *et al.* Clinicopathologic and genetic characterization of traditional serrated adenomas of the colon. *Am J Clin Pathol* 2012;138:356–66.
44. Tsai J-H, Liao J-Y, Lin Y-L, *et al.* Traditional serrated adenoma has two pathways of neoplastic progression that are distinct from the sessile serrated pathway of colorectal carcinogenesis. *Mod Pathol* 2014;27:1375–85.
45. East JE, Atkin WS, Bateman AC, *et al.* British Society of gastroenterology position statement on serrated polyps in the colon and rectum. *Gut* 2017;66:1181–96.
46. Gonzalo DH, Lai KK, Shadrach B, *et al.* Gene expression profiling of serrated polyps identifies annexin A10 as a marker of a sessile serrated adenoma/polyp. *J Pathol* 2013;230:420–9.
47. Kim JH, Rhee Y-Y, Kim K-J, *et al.* Annexin A10 expression correlates with serrated pathway features in colorectal carcinoma with microsatellite instability. *APMIS* 2014;122:1187–95.
48. Tsai J-H, Lin Y-L, Cheng Y-C, *et al.* Aberrant expression of annexin A10 is closely related to gastric phenotype in serrated pathway to colorectal carcinoma. *Mod Pathol* 2015;28:268–78.
49. Sajanti SA, Väyrynen JP, Simiö P, *et al.* Annexin A10 is a marker for the serrated pathway of colorectal carcinoma. *Virchows Arch* 2015;466:5–12.
50. Bae JMet *et al.* Annexin A10 expression in colorectal cancers with emphasis on the serrated neoplasia pathway. *WJG* 2015;21:9749–57.

51. Macaron C, Lopez R, Pai RK, *et al.* Expression of annexin A10 in serrated polyps predicts the development of metachronous serrated polyps. *Clin Transl Gastroenterol* 2016;7:e205.
52. Cui M, Awadallah A, Liu W, *et al.* Loss of HES1 differentiates sessile serrated adenoma/polyp from hyperplastic polyp. *Am J Surg Pathol* 2016;40:113–9.
53. Aoki H, Yamamoto E, Takasawa A, *et al.* Epigenetic silencing of SMOC1 in traditional serrated adenoma and colorectal cancer. *Oncotarget* 2017;9:4707–21.
54. Walsh MD, Buchanan DD, Pearson S-A, *et al.* Immunohistochemical testing of conventional adenomas for loss of expression of mismatch repair proteins in Lynch syndrome mutation carriers: a case series from the Australasian site of the colon cancer family registry. *Mod Pathol* 2012;25:722–30.
55. Jass JR. Classification of colorectal cancer based on correlation of clinical, morphological and molecular features. *Histopathology* 2007;50:113–30.
56. Bird AP. CpG-Rich islands and the function of DNA methylation. *Nature* 1986;321:209–13.
57. Toyota M, Ahuja N, Ohe-Toyota M, *et al.* CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci U S A* 1999;96:8681–6.
58. Yamane T *et al.* Serrated pathway in colorectal carcinogenesis. *WJG* 2014;20:2634–40.
59. Kim YH, Kakar S, Cun L, *et al.* Distinct CpG island methylation profiles and BRAF mutation status in serrated and adenomatous colorectal polyps. *Int J Cancer* 2008;23:2587–93.
60. Yang S, Farraye FA, Mack C, *et al.* Braf and KRAS mutations in hyperplastic polyps and serrated adenomas of the colorectum: relationship to histology and CpG island methylation status. *Am J Surg Pathol* 2004;28:1452–9.
61. Jass JR, Baker K, Zlobec I, *et al.* Advanced colorectal polyps with the molecular and morphological features of serrated polyps and adenomas: concept of a 'fusion' pathway to colorectal cancer. *Histopathology* 2006;49:121–31.
62. O'Brien MJ, Yang S, Mack C, *et al.* Comparison of microsatellite instability, CpG island methylation phenotype, BRAF and KRAS status in serrated polyps and traditional adenomas indicates separate pathways to distinct colorectal carcinoma end points. *Am J Surg Pathol* 2006;30:1491–501.
63. Sekine S, Yamashita S, Tanabe T, *et al.* Frequent *PTPRK-RSPO3* fusions and *RNF43* mutations in colorectal traditional serrated adenoma. *J Pathol* 2016;239:133–8.
64. Tsai J-H, Liao J-Y, Yuan C-T, *et al.* RNF43 is an early and specific mutated gene in the serrated Pathway, with increased frequency in traditional serrated adenoma and its associated malignancy. *Am J Surg Pathol* 2016;40:1352–9.
65. de Lau W, Peng WC, Gros P, *et al.* The R-spondin/Lgr5/Rnf43 module: regulator of Wnt signal strength. *Genes Dev* 2014;28:305–16.
66. Sekine S, Ogawa R, Hashimoto T, *et al.* Comprehensive characterization of *RSPO* fusions in colorectal traditional serrated adenomas. *Histopathology* 2017;71:601–9.
67. Yan HHN, Lai JCW, Ho SL, *et al.* RNF43 germline and somatic mutation in serrated neoplasia pathway and its association with BRAF mutation. *Gut* 2017;66:1645–56.
68. Gala MK, Mizukami Y, Le LP, *et al.* Germline mutations in oncogene-induced senescence pathways are associated with multiple sessile serrated adenomas. *Gastroenterology* 2014;146:520–9.
69. Taupin D, Lam W, Rangiah D, *et al.* A deleterious RNF43 germline mutation in a severely affected serrated polyposis kindred. *Hum Genome Var* 2015;2.
70. Bond CE, McKeone DM, Kalimutho M, *et al.* RNF43 and ZNRF3 are commonly altered in serrated pathway colorectal tumorigenesis. *Oncotarget* 2016;7:70589–600.
71. Liu C, McKeone DM, Walker NI, *et al.* GNAS mutations are present in colorectal traditional serrated adenomas, serrated tubulovillous adenomas and serrated adenocarcinomas with adverse prognostic features. *Histopathology* 2017;70:1079–88.
72. Yamada M, Sekine S, Ogawa R, *et al.* Frequent activating Gnas mutations in villous adenoma of the colorectum. *J Pathol* 2012;228:113–8.
73. Davis H, Irshad S, Bansal M, *et al.* Aberrant epithelial GREM1 expression initiates colonic tumorigenesis from cells outside the stem cell niche. *Nat Med* 2015;21:62–70.
74. Vogelstein B, Fearon ER, Hamilton SR, *et al.* Genetic alterations during colorectal-tumor development. *N Engl J Med* 1988;319:525–32.
75. Dubé C, McCurdy BR, Bronstein T, *et al.* *ColonCancerCheck recommendations for Post-Polypectomy surveillance*. Cancer Care Ontario, 2019.
76. Lieberman DA, Rex DK, Winawer SJ, *et al.* Guidelines for colonoscopy surveillance after screening and polypectomy: a consensus update by the US Multi-Society Task force on colorectal cancer. *Gastroenterology* 2012;143:844–57.
77. Leddin D, Enns R, Hilsden R, *et al.* Colorectal cancer surveillance after index colonoscopy: guidance from the Canadian association of gastroenterology. *Can J Gastroenterol* 2013;27:224–8.
78. Hassan C, Quintero E, Dumonceau J-M, *et al.* Post-polypectomy colonoscopy surveillance: European Society of gastrointestinal endoscopy (ESGE) guideline. *Endoscopy* 2013;45:842–64.
79. Snover DC. Diagnostic and reporting issues of preneoplastic polyps of the large intestine with early carcinoma. *Ann Diagn Pathol* 2019;39:1–14.