

Iron deficiency and symptoms in women aged 20–49 years and relation to upper gastrointestinal and colon cancers

Jean-Luc Szpakowski ,¹ Lue-Yen Tucker ²

To cite: Szpakowski J-L, Tucker L-Y. Iron deficiency and symptoms in women aged 20–49 years and relation to upper gastrointestinal and colon cancers. *BMJ Open Gastro* 2022;**9**:e000947. doi:10.1136/bmjgast-2022-000947

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgast-2022-000947>).

Received 3 May 2022
Accepted 6 July 2022

ABSTRACT

Objective Iron deficiency anaemia (IDA) in women aged 20–49 years may be caused by menses or gastrointestinal cancer. Data are sparse on the yield of endoscopy/colonoscopy in this population. Our aim was to determine the association of IDA and symptoms with cancers.

Design Retrospective cohort study within Kaiser Permanente Northern California. Participants were women aged 20–49 years tested for iron stores and anaemia during 1998, 2004 and 2010 and followed for 5 years for outcomes of oesophageal, gastric and colon cancers. Symptoms from the three prior years were grouped into dysphagia, upper gastrointestinal (UGI), lower gastrointestinal (LGI), rectal bleeding and weight loss.

Results Among 9783 anaemic women aged 20–49 years, there were no oesophageal, 6 gastric and 26 colon cancers. Incidences per 1000 for gastric cancer with and without iron deficiency (ID) were 0.60 (95% CI 0.23 to 1.55) and 0.63 (95% CI 0.17 to 2.31), and for colon cancer, 2.72 (95% CI 1.72 to 4.29) and 2.53 (95% CI 1.29 to 4.99). Endoscopies for UGI or dysphagia symptoms rather than bidirectional endoscopy for ID yielded more gastric cancers (n=5 and n=4, respectively) with fewer procedures (3793 instead of 6627). Colonoscopies for LGI or rectal bleed instead of for ID would detect more colon cancers (n=19 and n=18) with about 40% of the procedures (=2793/6627).

Conclusions UGI and colon cancers were rare in women of menstruating age and when controlled for anaemia were as common without as with ID. Using symptoms rather than IDA as an indication for endoscopy found equal numbers of cancers with fewer procedures.

INTRODUCTION

Iron deficiency anaemia (IDA) in adults in the developed world is commonly caused by bleeding in the gastrointestinal (GI) tract, with the most feared cause being cancer. The finding of IDA thus commonly leads to referral for endoscopic examination of the upper gastrointestinal (UGI) and lower gastrointestinal (LGI) tracts. In women of menstruating age, IDA is more commonly caused by blood loss with menses, occurring in 5%–15.7% of US women aged 20–49 years.^{1,2} This raises the question of whether these women should be offered endoscopy,

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Iron deficiency anaemia can be caused by gastrointestinal cancer, but in women of menstruating age, it is most commonly caused by menses. Professional organisations disagree whether their rate of cancer is sufficient to warrant endoscopic gastrointestinal examination irrespective of symptoms.

WHAT THIS STUDY ADDS

⇒ In anaemic women aged 20–49 years, gastrointestinal cancers were as common with iron deficiency as in those without iron deficiency. Performing endoscopies or colonoscopies for symptoms found more cancers with fewer procedures than performing endoscopy and colonoscopy in all those with iron deficiency.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ This study suggests that in anaemic women aged 20–49 years, the decision to perform UGI endoscopy or colonoscopy should be dictated by symptoms and not by iron deficiency.

given that GI cancers occasionally occur in this age group. Most studies of the incidence of malignancy in this population are small and performed in those referred for endoscopy, with case series ranging in size from 19 to 187 women.^{3–10} A recent Veterans Administration study was large and population based but did not examine the role of symptoms.¹¹ Data are even more scarce in those with iron deficiency (ID) without anaemia.¹² In order to examine the inter-relation of ID, anaemia and symptoms on the incidence of major GI cancers in women of menstruating age, we undertook a retrospective cohort study.

METHODS

We identified women between the ages of 20 and 49 years who had test results for ID and anaemia within Kaiser Permanente Northern California (KPNC) electronic health records (EHRs). The Kaiser population is



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Gastroenterology, Kaiser Permanente Northern California, Fremont, California, USA

²Division of Research, Kaiser Permanente Northern California, Oakland, California, USA

Correspondence to

Dr Jean-Luc Szpakowski;
jeanlucszp@gmail.com



representative of the overall Northern California population.¹³ Additional selection criteria included: (1) continuous membership in the study entry (or index) year; (2) continuous membership for 36 months preceding the index year without any endoscopic procedures in that time; and (3) 36 or more active membership months in 4 years following the index year or until an eligible cancer was diagnosed. Exclusion criteria included prior hysterectomy or diagnosis of gastrointestinal cancer or inflammatory bowel disease (IBD) and childbirth in index or preceding year. Entry years were 6 years apart, 1998, 2004 and 2010, with 1995 the first year for which a complete electronic data set was available. Each woman could be a study subject of one entry year only. The study entry (index) date was defined as the first date within a study year that tests for ID were done.

ID was defined as ferritin less than the lower limits of normal (LLN; 10–22 µg/L during this period). In those without available ferritins, we used total iron binding capacity greater than upper limits of normal (428 µg/dL), which has a higher correlation with ferritin than does iron saturation.^{12,13} Anaemia was defined as the LLN in our laboratories, haemoglobin <11.5 g/dL or haematocrit <34%, obtained during the year prior to the study entry date. When multiple tests were found, the test closest to the index date was used.

Women with anaemia were the primary cohort, and those without anaemia were a secondary cohort. Each cohort was further divided into: (1) ID versus no ID and (2) had endoscopy versus no endoscopy. We use 'endoscopy' in this study as a generic term encompassing any UGI endoscopy or colonoscopy performed within 1 year after the index date.

Outcomes

Primary outcomes included oesophageal cancers, gastric cancers and colon cancers within 5 years of the study entry date and were compiled from the KPNC Cancer Registry, with diagnosis confirmed by review of pathology records. Small carcinoids (<1 cm) were excluded as their biological behaviour is usually benign. We included two secondary outcomes: (1) we analysed findings in those undergoing endoscopy within 1 year of study entry date in order to examine biases introduced by selection for early endoscopy. (2) We examined the rates of selected non-cancer causes of IDA, acknowledging that their diagnosis is imperfect without endoscopy. The included causes were oesophagitis, ulcers or IBD, compiled from International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes.

Comorbidities and symptoms

Comorbidities and symptoms in the 3 years preceding the index date were identified using ICD-9-CM codes. Comorbidities of interest included diabetes mellitus, kidney disease, cardiovascular disease, pulmonary disease, menorrhagia and cirrhosis. Charlson

Comorbidity Index score was calculated using EHR records from this time period. Symptoms were grouped into five categories: (1) UGI symptoms, (2) dysphagia, (3) lower GI symptoms, (4) rectal bleeding and (5) weight loss (online supplemental table 1).

Patient demographics

Age at study entry date was calculated as a continuous variable and further grouped into 20–29, 30–39 and 40–49 years. Race/ethnicity included Hispanic/Latino, non-Hispanic white, non-Hispanic black, Asian/Pacific Islanders and other.

Statistical analysis

Distributions of demographics and clinical characteristics were calculated and expressed as frequencies (%) or mean (±SD) as appropriate. The incidence (per 1000 patients) of each cancer or secondary outcome was calculated by age groups, including 95% CI, according to: (1) ID status, (2) undergoing endoscopy within 1 year of ID testing or (3) presence or absence of symptom groups. We calculated the number needed to detect (NND) a cancer as a proxy for the number of endoscopies needed to detect one cancer and used it to compare the outcomes of different criteria for performing endoscopy. Comparisons in bivariate analyses included χ^2 or Fisher's exact tests for categorical variables as appropriate and independent two-sample t-tests for continuous variables.

Multivariable logistic regression analyses were used to assess the risk of being ID and (1) having cancer and (2) undergoing endoscopy, each stratified by anaemia status and controlling for symptom groups, patient race/ethnicity, Charlson score and patient age by deciles. All analyses were performed using SAS V.9.4 (SAS Institute Inc, Cary, North Carolina, USA) with the threshold of significance set at two-sided $p < 0.05$.

RESULTS

Findings in anaemic women

Demographics

A total of 32 440 women were tested for ID, of whom 9783 were anaemic (primary cohort), 22 657 were not anaemic (secondary cohort) and 942 had unknown status. The demographic and clinical characteristics of the anaemic cohort are shown in [table 1](#). Women with ID were more likely to be Hispanic and less likely to be black and had a lower Charlson score and more menorrhagia than non-ID anaemic women. Those who underwent endoscopy were more likely to be ID, older, white and had higher Charlson score and more symptoms.

Cancers

Thirty-two cancers were identified in anaemic women after we excluded one carcinoid less than 1 cm. There were no oesophageal, 6 gastric and 26 colon cancers.

Table 1 Demographic and clinical characteristics in women aged 20–49 years with anaemia who were tested for iron deficiency

Characteristics	Entire cohort n=9783 (100)	Iron deficiency n (%)		Had endoscopy* n (%)	
		Yes (n=6627)	No (n=3156)	Yes (n=653)	No (n=9130)
Iron deficiency	7919 (65.5)	–	–	527 (80.70%)	6100 (66.81%)
Had endoscopy*	1036 (8.6)	527 (7.95)	126 (3.99)	–	–
Age groups (year)					
20–29	648 (5.4)	379 (5.72)	269 (8.52)	15 (2.30)	633 (6.93)
30–39	2591 (21.4)	1656 (24.99)	935 (29.63)	118 (18.07)	2473 (27.09)
40–49	6544 (54.2)	4592 (69.29)	1952 (61.85)	520 (79.63)	6024 (65.98)
Age (year), mean±SD	43.1±7.3				
Charlson comorbidity index score		0.5±0.9	0.9±1.4	0.9±1.5	0.6±1.1
Mean±SD	0.7±1.2	2134 (32.2)	1367 (43.31)	282 (43.19)	3219 (35.26)
≥1	4568 (37.8)	379 (5.72)	269 (8.52)	15 (2.30)	633 (6.93)
Race/ethnicity					
White	3686 (30.5)	1886 (28.46)	865 (27.41)	228 (34.92)	2523 (27.63)
Black	2397 (19.8)	1178 (17.78)	793 (25.13)	95 (14.55)	1876 (20.55)
Hispanic	2480 (7.8)	1660 (25.05)	466 (14.77)	164 (25.11)	1962 (21.49)
Asian	2574 (21.3)	1417 (21.38)	754 (23.89)	114 (17.46)	2057 (22.53)
Other†	947 (7.8)	486 (7.33)	278 (8.81)	52 (7.96)	712 (7.80)
Any comorbidity	3292 (27.2)	1661 (25.06)	826 (26.17)	172 (26.34)	2315 (25.36)
Diabetes mellitus	1304 (10.8)	548 (8.27)	376 (11.91)	72 (11.03)	852 (9.33)
Kidney disease	333 (2.8)	66 (1.00)	162 (5.13)	21 (3.22)	207 (2.27)
Cardiovascular disease	238 (2.0)	63 (0.95)	72 (2.28)	13 (1.99)	12 (1.34)
Pulmonary disease	330 (2.7)	140 (2.11)	115 (3.64)	17 (2.60)	238 (2.61)
Menorrhagia	1712 (14.2)	1059 (15.98)	286 (9.06)	87 (13.32)	1258 (13.78)
Cirrhosis	40 (0.3)	10 (0.15)	17 (0.54)	9 (1.38)	18 (0.20)
Any symptoms	6372 (52.7)	3411 (51.47)	1667 (52.82)	506 (77.49)	4572 (50.08)
UGI symptoms	4688 (38.8)	2504 (37.78)	1201 (38.05)	424 (64.93)	3281 (35.94)
Dysphagia	390 (3.2)	183 (2.76)	103 (3.26)	44 (6.74)	242 (2.65)
LGI symptoms	3060 (25.3)	1622 (24.48)	849 (26.90)	251 (38.44)	2220 (24.32)
Rectal bleed	710 (5.9)	402 (6.07)	175 (5.54)	82 (12.56)	495 (5.42)
Weight loss	291 (2.4)	135 (2.04)	85 (2.69)	28 (4.29)	192 (2.10)
Any gastric symptoms: UGI and/or dysphagia	3705 (37.9)	2504 (37.8)	1201 (38.1)	424 (64.9)	3281 (35.9)
Any colon symptoms: LGI and/or rectal bleed	2471 (25.3)	1622 (24.5)	849 (26.9)	251 (38.4)	2220 (24.3)
Endoscopy within 5 years of study entry	1498 (15.3)	1101 (16.6)	397 (12.6)	653 (100)	845 (9.3)

Data are column % unless otherwise stated.

–, not applicable.

*Defined as undergoing endoscopy within 1 year of the iron deficiency test date.

†Including American Indians, Alaskan Natives, more than one racial groups and unknown.

LGI, lower gastrointestinal; UGI, upper gastrointestinal.

Gastric cancers were rare in all age cohorts, whereas colon cancer incidences increased with age (table 2). The overall incidence of GI malignancy was 3.32 cases per 1000 for those with ID and 3.17 for those without ID. The incidence of gastric cancer was similar with and without ID (0.60 (95% CI 0.23 to 1.55) and 0.63

(95% CI 0.17 to 2.31), respectively, $p>0.99$) as was the incidence of colon cancer (2.72 (95% CI 1.72 to 4.29) and 2.53 (95% CI 1.29 to 4.99), respectively, $p=0.81$).

When we examined those who had endoscopy in the first year, there was also no difference in cancer incidence in those with and without ID (table 3).



Table 2 Unadjusted incidence of gastric and colon cancers in women who were iron deficient compared with those who were not iron deficient

	Iron deficient			Non-iron deficient			P value
	Sample size (n)	Cancer* (n)	Incidence/1000 patients (95% CI)	Sample size (n)	Cancer* (n)	Incidence/1000 patients (95% CI)	
Anaemic women (n=9783)							
Gastric cancer							
Age 20–49 years	6627	4	0.60 (0.23 to 1.55)	3156	2	0.63 (0.17 to 2.31)	>0.99
20–29	379	0	–	269	0	–	–
30–39	1656	3	1.81 (0.62 to 5.31)	935	0	–	0.56
40–49	4592	1	0.22 (0.04 to 1.23)	1952	2	1.02 (0.28 to 3.73)	0.21
Colon cancer							
Age 20–49 years	6627	18	2.72 (1.72 to 4.29)	3156	8	2.53 (1.29 to 4.99)	0.87
20–29	379	1	2.64 (0.47 to 14.79)	269	0	–	>0.99
30–39	1656	5	3.02 (1.29 to 7.05)	935	2	2.14 (0.59 to 7.77)	>0.99
40–49	4592	12	2.61 (1.50 to 4.56)	1952	6	3.07 (1.41 to 6.69)	0.74
Non-anaemic women (n=22657)							
Gastric cancer							
Age 20–49 years	6979	0	–	15678	3	0.19 (0.07 to 0.56)	0.56
20–29	1010	0	–	2122	0	–	–
30–39	2068	0	–	5076	0	–	–
40–49	3901	0	–	8480	3	0.35 (0.12 to 1.04)	0.56
Colon cancer							
Age 20–49 years	6979	16	2.29 (1.41 to 3.72)	15678	12	0.77 (0.44 to 1.34)	0.003
20–29	1010	0	–	2122	0	–	–
30–39	2068	1	0.48 (0.09 to 2.73)	5076	1	0.20 (0.03 to 1.12)	0.50
40–49	3901	15	3.85 (2.33 to 6.33)	8480	11	1.30 (0.72 to 2.32)	0.01

Comparisons were made using Fisher's exact tests.

–, not applicable.

*Within 5 years of the iron deficiency test date.

Symptoms and cancer

Dysphagia was present in 286 women, but no oesophageal cancers were diagnosed. Gastric cancer had significant associations with UGI symptoms, dysphagia and weight loss. Colon cancer was associated with rectal bleeding and LGI and UGI symptoms. (table 4). Notably, ID was not associated with any primary outcome.

In multivariable modelling, the adjusted odds for gastric cancer were significantly increased with dysphagia (adjusted OR (aOR) 6.41, 95% CI 1.69 to 24.27) and weight loss (aOR 24.38, 95% CI 7.05 to 84.31), while the odds for colon cancer were increased with LGI (aOR 3.02, 95% CI 1.44 to 6.36) or UGI (aOR 2.49, 95% CI 1.15 to 5.39) symptoms and rectal bleeding (aOR 7.15, 95% CI 3.44 to 14.85) (online supplemental table 2). Of note, ID did not increase the odds of either cancer.

Odds of endoscopy

The adjusted odds of receiving endoscopy decreased with younger age and Asian, black or Hispanic race/ethnicity (online supplemental table 3). The odds increased with ID, any symptoms and higher Charlson score. On examining the interaction of symptoms and ID with receiving endoscopy, the ID women who received endoscopy had fewer LGI symptoms than did the non-ID women who received endoscopy (34.9% compared with 53.2%, $p < 0.001$), but the UGI/dysphagia symptoms were equal in the two groups (online supplemental table 4).

Optimal endoscopy strategies

The numbers of endoscopies needed to detect one cancer according to ID status and symptoms are presented in table 4. Endoscopy for ID had similar yield to endoscopy without ID. A strategy of UGI endoscopies performed only for UGI or dysphagia symptoms would yield more

Table 3 Unadjusted incidence of gastric and colon cancer in women who underwent endoscopy procedure within 1 year of testing for iron deficiency*

	Iron deficient			Non-iron deficient			P value
	Sample size (n)	Cancer† (n)	Incidence/1000 patients (95% CI)	Sample size (n)	Cancer† (n)	Incidence/1000 patients (95% CI)	
Anaemic women (n=653)							
Gastric cancer							
Age 20–49 years	527	1	1.90 (0.34 to 10.67)	126	1	7.94 (1.40 to 43.59)	0.35
20–29	11	0	–	4	0	–	–
30–39	82	1	12.20 (2.16 to 65.89)	36	0	–	>0.99
40–49	434	0	–	86	1	11.63 (2.06 to 62.96)	0.17
Colon cancer							
Age 20–49 years	527	8	15.18 (7.71 to 29.67)	126	2	15.87 (4.36 to 56.03)	>0.99
20–29	11	0	–	4	0	–	–
30–39	82	4	48.78 (19.13 to 118.80)	36	1	27.78 (4.92 to 141.70)	>0.99
40–49	434	4	9.22 (3.59 to 23.46)	86	1	11.63 (2.06 to 62.96)	>0.99
Non†anaemic women (n=939)							
Gastric cancer							
Age 20–49 years	388	0	–	551	0	–	–
20–29	33	0	–	47	0	–	–
30–39	75	0	–	132	0	–	–
40–49	280	0	–	372	0	–	–
Colon cancer							
Age 20–49 years	388	7	18.04 (8.77 to 36.77)	551	3	5.44 (1.85 to 15.88)	0.10
20–29	33	0	–	47	0	–	–
30–39	75	1	13.33 (2.36 to 71.73)	132	1	7.58 (1.34 to 41.66)	>0.99
40–49	280	6	21.43 (9.86 to 45.95)	372	2	5.38 (1.48 to 19.39)	0.08

–, not applicable.
 *No comparisons were significant due to small event size.
 †Within 5 years of the iron deficiency test date.

gastric cancers than using ID as a criterion (n=5 and n=4, respectively), with 57.2% (=3793/6627) the number of UGI endoscopies. Performing colonoscopies only for LGI or rectal bleed symptoms instead of for ID would yield more colon cancers (n=19 and n=18, respectively), with 42.1% (=2793/6627) the number of colonoscopies. A detailed summary of the NND according to symptom and age subgroups is available in online supplemental table 5.

A key question is that of yield of endoscopy in women with IDA but without gastrointestinal symptoms. In those with IDA without dysphagia or UGI symptoms, no gastric cancers were found in 4064 women aged 20–49 years. In those with IDA but without LGI symptoms or rectal bleeding, four colon cancers were found in 4778 women for a rate of 0.84 per 1000, far less than the rate of 2.66 per 1000 in the entire cohort.

Non-cancer findings

The numbers of oesophagitis, ulcers and IBD diagnosed both clinically and endoscopically over the subsequent 5 years is shown in table 5, and the numbers diagnosed

by endoscopy within the first year are shown in online supplemental table 6. There was no increase in these diagnoses with ID; if anything, the trend was towards more findings in those without ID. Because oesophagitis and ulcers would be diagnosed by UGI endoscopy, we investigated whether this trend was because a higher percentage of non-ID underwent UGI endoscopy for UGI symptoms, but we found that the rate of UGI symptoms was similar in those with and without ID (online supplemental table 4).

Findings in non-anaemic women

The demographics of the women tested for ID and without anaemia were similar to those with anaemia (online supplemental table 7). Of note, those with ID had more menorrhagia, and those who underwent endoscopy had more symptoms and a higher Charlson score.

The overall findings were similar to those of anaemic patients, although with generally lower rates of cancer and therefore wider CIs (table 2). The presence or absence of ID made no difference in the incidence of cancers, with the exception of more colon cancers in



Table 4 Unadjusted incidence of gastric and colon cancers stratified by iron deficiency and symptoms in 9783 anaemic women aged 20–49 years

	Sample size (n)	Cancer* (n)	NND one cancer (95% CI)	Incidence/1000 patients (95% CI)	P value
Gastric cancer, age 20–49 years					
Random	9783	6	1631 (748 to 3558)	0.61 (0.28 to 1.34)	–
Iron deficiency					
(+)	6627	4	1657 (645 to 4260)	0.60 (0.23 to 1.55)	>0.99
(–)	3156	2	1578 (434 to 5754)	0.63 (0.17 to 2.31)	
Any symptoms					
Yes	5078	6	847 (389 to 1847)	1.18 (0.54 to 2.58)	0.03
No	4705	0	–	–	
UGI symptoms					
Yes	3705	5	741 (317 to 1735)	1.35 (0.58 to 3.16)	0.03
No	6078	1	6078 (1074 to 34,431)	0.16 (0.03 to 0.93)	
Dysphagia					
Yes	286	2	143 (40 to 521)	6.99 (1.92 to 25.10)	0.01
No	9497	4	2375 (924 to 6105)	0.42 (0.16 to 1.08)	
Weight loss					
Yes	220	3	74 (26 to 216)	13.6 (4.65 to 39.3)	<0.001
No	9563	3	3188 (1085 to 9373)	0.31 (0.11 to 0.92)	
Any gastric symptoms: UGI and/or dysphagia					
All women					
Yes	3793	5	759 (325 to 1776)	1.32 (0.56 to 3.08)	0.04
No	5990	1	5990 (1059 to 33,933)	0.17 (0.03 to 0.95)	
Iron deficient (+) women (n=6627)					
Yes	2563	4	641 (250 to 1648)	1.56 (0.61 to 4.01)	0.02
No	4064	0	–	–	
Iron deficient (–) women (n=3156)					
Yes	1230	1	1230 (218 to 6968)	0.81 (0.14 to 4.59)	>0.99
No	1926	1	1926 (341 to 10,910)	0.52 (0.09 to 2.94)	
Colon cancer, age 20–49 years					
Random	9783	26	377 (257 to 552)	2.66 (1.81 to 3.89)	–
Iron deficiency					
(+)	6627	18	369 (234 to 582)	2.72 (1.72 to 4.29)	>0.99
(–)	3156	8	395 (201 to 779)	2.53 (1.29 to 4.99)	
Any symptoms					
Yes	5078	21	242 (159 to 370)	4.14 (2.71 to 6.31)	0.003
No	4705	5	941 (403 to 2203)	1.28 (0.45 to 2.49)	
LGI symptoms					
Yes	2471	16	155 (96 to 251)	6.48 (3.99 to 10.50)	<0.001
No	7312	10	732 (398 to 1346)	1.37 (0.74 to 2.52)	
Rectal bleed					
Yes	577	10	58 (32 to 106)	17.30 (9.44 to 31.60)	<0.001
No	9206	16	576 (355 to 935)	1.74 (1.07 to 2.82)	
Weight loss					
Yes	220	1	220 (40 to 1246)	4.55 (0.80 to 25.30)	0.48
No	9563	25	383 (260 to 565)	2.61 (1.77 to 3.86)	

Continued

**Table 4** Continued

	Sample size (n)	Cancer* (n)	NND one cancer (95% CI)	Incidence/1000 patients (95% CI)	P value
Any colon symptoms: LGI and/or rectal bleed					
All women					
Yes	2793	19	147 (95 to 230)	6.80 (4.36 to 10.60)	<0.001
No	6990	7	999 (485 to 2062)	1.00 (0.49 to 2.07)	
Iron deficient (+) women (n=6627)					
Yes	1849	14	133 (79 to 222)	7.57 (4.52 to 12.70)	<0.001
No	4778	4	1195 (465 to 3072)	0.84 (0.33 to 2.15)	
Iron deficient (-) women (n=3156)					
Yes	944	5	189 (82 to 442)	5.30 (2.26 to 12.30)	0.02
No	2212	3	738 (252 to 2168)	1.36 (0.46 to 3.98)	

Comparisons were made using Fisher's exact tests.

*Within 5 years of the iron deficiency test date.

(-), negative; (+), positive; -, not applicable; LGI, lower gastrointestinal; NND, number needed to detect; UGI, upper gastrointestinal.

women aged 40–49 years with ID, which in turn led to significance in the differences in the combined 20–49 cohort. This 40–49 ID cohort was a further anomaly as the only age bracket where the incidence of cancer was higher in non-anaemic women than in anaemic women. In the subgroup with endoscopy, ID made no difference in cancer incidence (table 3).

The associations with specified symptoms were of lower frequencies than in anaemic women but again using symptoms rather than ID as a guide to endoscopy found equivalent number of cancers with fewer endoscopies (online supplemental tables 8). Notably, in women without UGI or dysphagia symptoms, no gastric cancers were found in 4178 women with ID nor in 9511 without ID. In those without rectal bleed or LGI symptoms, six colon cancers were found in 4747 women with ID for a rate of 1.26 per 1000, about the same rate as colon cancer in a random woman (1.24 per 1000) and below the rate of 2.03 in those with symptoms regardless of ID status.

For non-cancer findings, again, there was no significant difference with or without ID in those without anaemia (table 5).

DISCUSSION

Our study found that in women aged 20–49 years who were tested for ID and had anaemia, the incidence of gastrointestinal cancers was low and was not affected by the presence or absence of ID. Our incidences are lower than in older studies most likely because those study cohorts were small and included only patients who received endoscopy. Those studies are therefore comparable with our cohort who had IDA and endoscopy within a year. For example, when we summed the rates in eight prior case series in women aged 20–49 years, 8 out of 818 women (0.98%) had gastric cancers and 14 of 600 (2.33%) had colon cancer.^{3–10} In our study, the incidence in those who had IDA and early

endoscopy was 0.19% and 1.52%, respectively. When we included the entire population who tested positive for IDA regardless of endoscopy, our incidence was lower, 0.06% for gastric cancers and 0.30% for colon cancers. By comparison, this is similar to the general female population incidence of acquiring gastric cancer (0.1%) and colon cancer (0.4%) between the ages of 20 and 49 years.¹⁴

In reviewing the risk factors for early endoscopy, we showed a selection bias in that those who receive endoscopy were more likely to be ID and to have symptoms. This is important because those with ID but without symptoms had a much lower rate of cancer, lower even than the rate of cancer in anaemic women randomly endoscoped.

This suggests that primary care doctors and gastroenterologists were selective in referring patients and in performing endoscopies even within the IDA group.

We are aware of two other population studies in the 20–49 age group. A 2-year follow-up of the 1971–1974 National Health and Nutrition Examination Survey found that none of the 92 premenopausal women with IDA were diagnosed with gastrointestinal malignancy, nor were 350 ID without anaemia.¹⁵ Second, in 2020, a large Veterans Administration study was published examining the rate of colon cancer in those with IDA and age 18–49 years.¹¹ Their 5-year cumulative incidence in women aged 18–49 years was 0.11%, lower than in our study. Unlike our study, their controls were all those without IDA without requirement for iron testing nor stratifying by anaemia status, nor did they examine the role of symptoms.

In agreement with other studies and the American Gastroenterological Association (AGA) recommendations, we found that those with symptoms had a higher incidence of malignancy than those without symptoms.^{3,4,16} Indeed, we found that a strategy of UGI endoscopy in those with UGI or dysphagia symptoms

Table 5 Unadjusted incidence of oesophagitis, ulcers and IBD in women who were iron deficient compared with those who were not iron deficient

	Iron deficient			Non-iron deficient			P value
	Sample size (n)	Pathology* (n)	Incidence/1000 patients (95% CI)	Sample size (n)	Pathology* (n)	Incidence/1000 patients (95% CI)	
Anaemic women (n=9783)							
Oesophagitis							
Age 20–49 years	6627	100	15.09 (12.42 to 18.32)	3156	66	20.91 (16.47 to 26.52)	0.04
20–29	379	6	15.83 (7.28 to 34.10)	269	2	7.43 (2.04 to 26.70)	0.48
30–39	1656	28	16.91 (11.72 to 24.33)	935	24	25.67 (17.31 to 37.91)	0.13
40–49	4592	66	14.37 (11.31 to 18.24)	1952	40	20.49 (15.08 to 27.78)	0.07
Ulcer							
Age 20–49 years	6627	59	8.90 (6.91 to 11.47)	3156	41	12.99 (9.59 to 17.58)	0.06
20–29	379	1	2.64 (0.47 to 14.79)	269	3	11.15 (3.80 to 32.27)	0.31
30–39	1656	10	6.04 (3.28 to 11.08)	935	6	6.42 (2.94 to 13.93)	0.90
40–49	4592	48	10.45 (7.89 to 13.83)	1952	32	16.39 (11.64 to 23.05)	0.045
IBD							
Age 20–49 years	6627	24	3.62 (2.43 to 5.38)	3156	14	4.44 (2.64 to 7.43)	0.54
20–29	379	1	2.64 (0.47 to 14.79)	269	1	3.72 (0.66 to 20.75)	>0.99
30–39	1656	5	3.02 (1.29 to 7.05)	935	4	4.28 (1.66 to 10.95)	0.73
40–49	4592	18	3.92 (2.48 to 6.19)	1952	9	4.61 (2.43 to 8.74)	0.69
Non-anaemic women (n=22 657)							
Oesophagitis							
Age 20–49 years	6979	82	11.75 (9.48 to 14.56)	15678	221	14.10 (12.37 to 16.06)	0.16
20–29	1010	6	5.94 (2.73 to 12.90)	2122	17	8.01 (5.01 to 12.79)	0.53
30–39	2068	22	10.64 (7.04 to 16.06)	5076	56	11.03 (8.51 to 14.30)	0.88
40–49	3901	54	13.84 (10.63 to 18.02)	8480	148	17.45 (14.88 to 20.47)	0.14
Ulcer							
Age 20–49 years	6979	58	8.31 (6.43 to 10.73)	15678	97	6.19 (5.07 to 7.54)	0.07
20–29	1010	3	2.97 (1.01 to 8.70)	2122	8	3.77 (1.91 to 7.42)	>0.99
30–39	2068	17	8.22 (5.14 to 13.13)	5076	26	5.12 (3.50 to 7.49)	0.12
40–49	3901	38	9.74 (7.11 to 13.34)	8480	63	7.43 (5.81 to 9.49)	0.18
IBD							
Age 20–49 years	6979	30	4.30 (3.01 to 6.13)	15678	65	4.15 (3.25 to 5.28)	0.87

Continued

Table 5 Continued

	Iron deficient		Non-iron deficient		P value
	Sample size (n)	Pathology* (n)	Sample size (n)	Pathology* (n)	
20–29	1010	0	2122	7	0.10
30–39	2068	6	5076	15	0.97
40–49	3901	24	8480	43	0.45
					Incidence/1000 patients (95% CI)
					3.30 (1.60 to 6.79)
					2.96 (1.79 to 4.87)
					4.95 (3.67 to 6.69)

Comparisons were made using Fisher's exact tests.
 –, not applicable.
 *Within 5 years of the iron deficiency test date.
 IBD, inflammatory bowel disease.

and colonoscopy for those with rectal bleeding or LGI symptoms was overall superior to one of endoscopy for IDA in terms of yield of cancers and number of endoscopies needed to detect them. This calls into question the usefulness of routine bidirectional ('flip') procedures for ID done in order to discover asymptomatic cancers in these age groups. In particular, gastric cancers without UGI symptoms were extremely rare in this age group. In women without gastric symptoms, no gastric cancers were found in 4064 ID women with anaemia or 4178 without anaemia. If confirmed in other studies, this low yield would call into question routine UGI endoscopy for asymptomatic women in these age groups with ID and suggests instead targeting endoscopies at those with symptoms. For colon cancers, performing colonoscopies in asymptomatic women with ID is no more productive than targeting a random woman tested for ID. Equivalent number of cancers were found with far fewer procedures by using symptoms instead of ID as an indication for colonoscopy.

Our paper focuses on cancers, but we included the rates of selected non-cancer findings in order to give a rough estimate for those findings and to acknowledge that these findings may also be important and may justify endoscopy. Unlike cancers, these may remain asymptomatic during the follow-up period and thus their diagnosis without UGI endoscopy is less reliable. We did find that in those who underwent endoscopy in the first year the rate of findings did not differ by ID status. We reviewed whether this equivalence was due to a confounding by indication, whereby in non-ID women, more endoscopies were performed for symptoms, but we found that they had the same rate of UGI symptoms as did those with ID. This suggests that it is not a given that young women with ID have more of these findings, indeed the trend in our study was in the opposite direction. Our findings are hypothesis generating, as it would take a separate and larger study to fully account for all the confounders and biases in looking at this issue, including the role of asymptomatic findings in this age group.

Our findings support the British Society of Gastroenterology 2021 recommendation that per se 'IDA in young women is not an indication for endoscopic investigation' and should be reserved for the presence of red flag symptoms, major genetic risk, or recurrent or refractory anaemia disproportionate to other causes, or absence of menses.¹⁷ Our findings support the AGA technical review recommendation that in patient who have gastrointestinal symptoms, evaluation should be site directed rather than routine bidirectional endoscopy.¹⁶ Our paper does not support the 2020 AGA recommendation for bidirectional endoscopy over iron replacement for asymptomatic premenopausal women with IDA and agrees with its conclusion that the pooled evidence overestimates the incidence of cancer due to inclusion of symptomatic



women.¹⁸ We also found that symptoms were a more efficient indication for endoscopy than ID, as ID in the absence of symptoms did not increase the risk of cancer in women with anaemia. Finally, a subsequent published correspondence with the authors of the guideline noted that the AGA guideline did not address ID without anaemia in large part due to a 'paucity of data'.^{12 19} Our findings help address this deficit. Whether (fecal immunochemical test (FIT)) testing can be used to further stratify who in this age group should undergo endoscopy, as has been shown to be the case in older age groups, can be tested in future studies.²⁰

Strengths

Our study avoids referral bias by including all patients diagnosed with ID, not just those referred for endoscopic examination, and ensured adequate follow-up by its inclusion criteria.

Our definition of ID included a ferritin LLN as that report label highlighted a value that would trigger attention and action. What levels to choose as a cut-off depends on the values attached to sensitivity versus specificity and on the presence of 'inflammatory diseases',²¹ many of which are components of the Charlson score.²² In women of childbearing age, who are generally healthy,²³ a level <15 µg/L is recommended by the WHO and has been shown to have a specificity of 98% and sensitivity of 75% for ID as defined by no stainable bone marrow iron.^{23 24} Our level was close to this, and the low Charlson score in our patients is consistent with their being generally healthy.

Limitations

An inherent limitation of our study is that endoscopy is the method most used to diagnose gastric and colon cancers, raising the possibility of underdiagnosis in the group without early endoscopy. We addressed this by including all cancers diagnosed in the first 5 years, expecting that cancers large enough to cause ID would become apparent within 5 years. This is similar to assumptions in other studies although they used 2 years of follow-up.^{15 25} To control for the possibility that de novo cancers in the years after diagnosis of ID attenuated the difference between ID and non-ID, we examined the differences in those with endoscopy within a year, but here too found no difference.

We relied on symptoms as coded by physicians and we cannot rule out that uncoded diagnoses exist, despite the fact that a high percentage of our study group had symptoms. The symptom of rectal bleeding or haematochezia deserves special attention in formulating algorithms in light of population studies showing that 15%–18% of the population has some rectal bleeding every year.^{26 27} Thus, the specificity and workload will vary greatly depending on how that symptom is defined.

Our cohorts included only those tested for ID, which was necessary in order to derive a comparison group without ID. We do not address the risk of cancer in those not tested for ID nor do we address why patients were tested for ID, whether as a routine screen or as a targeted test. In this, we address the problem as faced by the gastroenterologist.

The number of outcome events was low in our study, with wide CIs, so the results must be interpreted with caution and confirmed by other studies. However, the rarity of outcome events in those with IDA is in itself an important finding and perhaps not unexpected by clinicians.²⁸ The even lower rate of findings in those with ID without anaemia can help guide decisions on whether endoscopy should be offered to these women given the controversy provoked by the AGA guidelines.

CONCLUSION

Our population-wide study found that UGI and colon cancers were rare in women of menstruating age and that when controlled for the presence of anaemia, they were not more common in those who were ID than in those who were not. We also found that in this age group, a symptom-based strategy detected more cancers with fewer procedures than did a strategy of endoscopy for those with ID.

Twitter Jean-Luc Szpakowski @Lucszpakowski

Contributors J-LS was responsible for initial conception and first draft, L-YT for initial data analysis, and both contributed to further design, analysis and interpretation of the data, critical revision of the article for important intellectual content, and both gave final approval of the article. J-LS is the guarantor of the overall content of this article.

Funding This work was supported by Kaiser Permanente Northern California Community Health grant number 14-2028-H.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study was approved by the Kaiser Foundation Research Institute's Institutional Review Board with waiver of consent. This work was supported by Kaiser Permanente Northern California Community Health grant number 14-2028-H.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

 Jean-Luc Szpakowski <http://orcid.org/0000-0001-5795-2334>

 Lue-Yen Tucker <http://orcid.org/0000-0001-7980-9272>

REFERENCES

- 1 Centers for Disease Control and Prevention (CDC). Iron deficiency—United States, 1999–2000. *MMWR Morb Mortal Wkly Rep* 2002;51:897–9.
- 2 Cogswell ME, Looker AC, Pfeiffer CM, *et al*. Assessment of iron deficiency in US preschool children and nonpregnant females of childbearing age: National health and nutrition examination survey 2003–2006. *Am J Clin Nutr* 2009;89:1334–42.
- 3 Vannella L, Aloe Spiriti MA, Cozza G, *et al*. Benefit of concomitant gastrointestinal and gynaecological evaluation in premenopausal women with iron deficiency anaemia. *Aliment Pharmacol Ther* 2008;28:422–30.
- 4 Green BT, Rockey DC. Gastrointestinal endoscopic evaluation of premenopausal women with iron deficiency anemia. *J Clin Gastroenterol* 2004;38:104–9.
- 5 Park DI, Ryu SH, Oh SJ, *et al*. Significance of endoscopy in asymptomatic premenopausal women with iron deficiency anemia. *Dig Dis Sci* 2006;51:2372–6.
- 6 Fireman Z, Zachlka R, Abu Mouch S, *et al*. The role of endoscopy in the evaluation of iron deficiency anemia in premenopausal women. *Isr Med Assoc J* 2006;8:88–90.
- 7 Kepczyk T, Cremins JE, Long BD, *et al*. A prospective, multidisciplinary evaluation of premenopausal women with iron-deficiency anemia. *Am J Gastroenterol* 1999;94:109–15.
- 8 Nahon S, Lahmek P, Lesgourgues B, *et al*. Predictive factors of GI lesions in 241 women with iron deficiency anemia. *Am J Gastroenterol* 2002;97:590–3.
- 9 Bini EJ, Micale PL, Weinshel EH. Evaluation of the gastrointestinal tract in premenopausal women with iron deficiency anemia. *Am J Med* 1998;105:281–6.
- 10 Annibale B, Lahner E, Chistolini A, *et al*. Endoscopic evaluation of the upper gastrointestinal tract is worthwhile in premenopausal women with iron-deficiency anaemia irrespective of menstrual flow. *Scand J Gastroenterol* 2003;38:239–45.
- 11 Demb J, Liu L, Murphy CC, *et al*. Young-Onset colorectal cancer risk among individuals with iron-deficiency anaemia and haematochezia. *Gut* 2020. doi:10.1136/gutjnl-2020-321849. [Epub ahead of print: 18 Dec 2020].
- 12 Hew S, Goss K. AGA clinical practice guidelines on the gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2021;160:2616.
- 13 Gordon NP. Similarity of adult Kaiser Permanente members to the adult population in Kaiser Permanente's Northern California service area: comparisons based on the 2017/2018 cycle of the California Health Interview Survey. Report prepared for the Kaiser Permanente Division of Research, Oakland, CA, 202a. Available: https://divisionofresearch.kaiserpermanente.org/projects/memberhealthsurvey/SiteCollectionDocuments/compare_kp_ncal_chis2017-18.pdf
- 14 Society AC. Lifetime risk or risk for those who are cancer free at the beginning of selected age interval. Data sources: DevCan version 6.7, National Cancer Institute, 2019]. 2019. Available: <https://cancerstatisticscenter.cancer.org/cancer-site/Stomach/RsPI71qd> [Accessed 07 Oct 2020].
- 15 Ioannou GN, Rockey DC, Bryson CL, *et al*. Iron deficiency and gastrointestinal malignancy: a population-based cohort study. *Am J Med* 2002;113:276–80.
- 16 Rockey DC, Altayar O, Falck-Ytter Y, *et al*. AGA technical review on gastrointestinal guidelines for the management of iron deficiency anemia. *Gastroenterology* 2020;159:1097–119.
- 17 Snook J, Bhala N, Beales ILP, *et al*. British Society of gastroenterology guidelines for the management of iron deficiency anaemia in adults. *Gut* 2021;70:2030–51.
- 18 Ko CW, Siddique SM, Patel A, *et al*. AGA clinical practice guidelines on the gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2020;159:1085–94.
- 19 Ko CW, Siddique SM, Patel A. Reply. *Gastroenterology* 2021;160:2620–1.
- 20 D'Souza N, Georgiou Delisle T, Chen M, *et al*. Faecal immunochemical test is superior to symptoms in predicting pathology in patients with suspected colorectal cancer symptoms referred on a 2WW pathway: a diagnostic accuracy study. *Gut* 2021;70:1130–8.
- 21 Guyatt GH, Oxman AD, Ali M, *et al*. Laboratory diagnosis of iron-deficiency anemia: an overview. *J Gen Intern Med* 1992;7:145–53.
- 22 Quan H, Sundararajan V, Halfon P, *et al*. Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005;43:1130–9.
- 23 Clucas D, Brittenham G, Pasricha S-R. AGA clinical practice guidelines on the gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2021;160:2618–20.
- 24 Hallberg L, Bengtsson C, Lapidus L, *et al*. Screening for iron deficiency: an analysis based on bone-marrow examinations and serum ferritin determinations in a population sample of women. *Br J Haematol* 1993;85:787–98.
- 25 Allison JE, Tekawa IS, Ransom LJ, *et al*. A comparison of fecal occult-blood tests for colorectal-cancer screening. *N Engl J Med* 1996;334:155–60.
- 26 Eslick GD, Kalantar JS, Talley NJ. Rectal bleeding: epidemiology, associated risk factors, and health care seeking behaviour: a population-based study. *Colorectal Dis* 2009;11:921–6.
- 27 Talley NJ, Jones M. Self-reported rectal bleeding in a United States community: prevalence, risk factors, and health care seeking. *Am J Gastroenterol* 1998;93:2179–83.
- 28 Munro MG, Critchley HOD, Lumsden MA. Re: Ko *et al*. AGA Clinical Practice Guidelines on the Gastrointestinal Evaluation of Iron Deficiency Anemia. *Gastroenterology* 2021;160:2616–7.

Iron deficiency and symptoms in women age 20-49 and relation to upper gastrointestinal and colon cancers: On-line supplement

eTable 1. Symptom categories and ICD-9-CM diagnosis codes.

eTable 2. Adjusted odds ratios for cancers in women age 20-49 with anemia who were tested for iron deficiency

eTable 3. Adjusted odd ratios of receiving endoscopy in women age 20-49 who were tested for iron deficiency.

eTable 4. Symptoms among women age 20-49 who underwent endoscopy within one year of testing for iron-deficiency

eTable 5. Sample number needed to detect a cancer in women with anemia age 20-49 with selected symptoms-by age subgroups.

eTable 6. Unadjusted incidence of esophagitis, ulcers and IBD in women who underwent endoscopy procedure within one year of testing for iron deficiency

eTable 7. Demographic and clinical characteristics in women age 20-49 without anemia who were tested for iron deficiency.

eTable 8. Unadjusted incidence of gastric and colon cancers stratified by iron deficiency and symptoms in 22,657 non-anemic women age 20-49.

eTable 1. Symptom categories and ICD-9-CM diagnosis codes

Category	Symptom	ICD-9-CM
Upper gastrointestinal symptoms		
	upper abdominal pain	789.01, 789.02
	epigastric pain	789.06
	dyspepsia	536.8
	gastritis	535
	gastrointestinal reflux	530.81
Dysphagia	dysphagia	438.82, 787.2
Lower gastrointestinal symptoms		
	change in bowel habits	787.99
	constipation	564
	diarrhea	787.91
	left lower quadrant pain	789.04
	right lower quadrant pain	789.03
	irritable bowel	564.1
Rectal bleed	rectal bleed	569.3
Weight loss	weight loss	783.21

eTable 2. Adjusted odds ratios for cancers in women age 20-49 with anemia who were tested for iron deficiency

Risk Factors	Age 20-49	
	Gastric Cancer ^a aOR (95% CI)	Colon Cancer ^a aOR (95% CI)
Iron deficiency	1.08 (0.32-3.65)	1.11 (0.52-2.37)
Any UGI symptoms	4.03 (0.98-16.66)	2.49 (1.15-5.39)
Any dysphagia symptoms	6.41 (1.69-24.27)	0.32 (0.03-4.25)
Any LGI symptoms	0.92 (0.26-3.28)	3.02 (1.44-6.36)
Any rectal bleed	1.12 (0.12-10.66)	7.15 (3.44-14.85)
Any weight loss	24.38 (7.05-84.31)	1.64 (0.35-7.68)
Age group ^b , years		
20-29	0.99 (0.08-12.88)	0.90 (0.19-4.22)
30-39	2.84 (0.87-9.31)	1.10 (0.50-2.43)
40-49 (reference group)	1	1
Race/Ethnicity		
White (reference group)	1	1
Asian	0.75 (0.13-4.45)	1.02 (0.42-2.48)
Black	0.73 (0.13-4.18)	0.42 (0.14-1.27)
Hispanic	1.31 (0.30-5.80)	0.42 (0.15-1.19)
Other ^c	0.87 (0.07-10.38)	0.42 (0.09-2.08)
Charlson comorbidity index score ≥1	1.04 (0.31-3.47)	2.00 (0.97-4.09)

Multivariable logistic regression analyses were used to assess the risk of being iron deficient and having cancer in women age 20-49 and 50-55, controlling for symptom groups, patient race/ethnicity, and Charlson score.

In bold are risk factors that are statistically significant at two-sided $P < 0.05$.

ID = iron deficiency; aOR = adjusted odds ratio; UGI = upper gastrointestinal; LGI = lower gastrointestinal; -- = not applicable.

^a Within five years of the iron deficiency test date.

^b Not included in the multivariable logistic regression modeling for colon cancer in women aged 50-55.

^c Including American Indians, Alaskan Natives, more than one racial groups, and unknown.

eTable 3. Adjusted odd ratios of receiving endoscopy^a in women age 20-49 who were tested for iron deficiency.

Risk Factors	Age 20-49	
	Anemic	Non-Anemic
	aOR (95% CI)	aOR (95% CI)
Iron Deficient	2.08 (1.69-2.55)	1.66 (1.45-1.91)
Any UGI symptom	2.82 (2.37-3.35)	2.79 (2.40-3.24)
Any dysphagia	1.77 (1.25-2.50)	2.02 (1.56-2.61)
Any LGI symptom	1.43 (1.20-1.70)	2.18 (1.89-2.51)
Any rectal bleed	1.97 (1.52-2.55)	2.97 (2.49-3.55)
Any weight loss	1.61 (1.06-2.46)	2.61 (1.99-3.43)
Age group ^b , years		
20-29	0.33 (0.19-0.55)	0.50 (0.39-0.63)
30-39	0.61 (0.49-0.75)	0.57 (0.49-0.68)
40-49 (reference group)	1	1
Race/Ethnicity		
White (reference group)	1	1
Asian	0.70 (0.55-0.89)	0.78 (0.63-0.96)
Black	0.60 (0.46-0.77)	0.65 (0.49-0.85)
Hispanic	0.79 (0.64-0.98)	0.82 (0.69-0.97)
Other ^c	0.79 (0.57-1.08)	0.90 (0.70-1.17)
Charlson comorbidity index score ≥ 1	1.22 (1.03-1.44)	1.24 (1.08-1.42)

Multivariable logistic regression analyses were used to assess the risk of having endoscopy within one year of testing for iron deficiency in women age 20-49 and 50-55, controlling for symptom groups, patient race/ethnicity, and Charlson score.

In bold are risk factors that are statistically significant.

ID = iron deficiency; aOR = adjusted odds ratio; UGI = upper gastrointestinal; LGI = lower gastrointestinal; -- = not applicable.

^a Defined as undergoing endoscopy within one year of the iron deficiency test date.

^b Not included in the multivariable logistic regression modeling for receiving endoscopy in women aged 50-55.

^c Including American Indians, Alaskan Natives, more than one racial groups, and unknown.

eTable 4. Symptoms among women age 20-49 who underwent endoscopy within one year of testing for iron-deficiency

Symptom	Iron Deficient n= 527)		Non-Iron Deficient n= 126		P value
Anemic					
Any	398	75.5%	108	85.7%	.01
UGI symptoms	336	63.8%	88	69.8%	.20
Dysphagia	35	6.6%	9	7.1%	.84
UGI and/or dysphagia	342	64.9%	91	72.2%	.12
LGI symptoms	184	34.9%	67	53.2%	<.001
Rectal bleed	67	12.7%	15	11.9%	.81
LGI and/or rectal bleed	307220	41.8%	75	59.5%	<.001
Non-anemic					
Any	342	88.1%	495	89.8%	.41
UGI symptoms	273	70.5%	386	70.1%	.92
Dysphagia	31	8.0%	52	9.4%	.44
UGI and/or dysphagia	277	71.4%	393	71.3%	.98
LGI symptoms	208	53.6%	320	58.1%	.17
Rectal bleed	72	18.6%	123	22.3%	.16
LGI and/or rectal bleed	235	60.6%	376	68.2%	.02

Comparisons were made using Fisher's exact tests.

eTable 5. Sample number needed to detect a cancer in women with anemia age 20-49 with selected symptoms-by age subgroups.

Sample size/n of cancers^a = number needed to detect one cancer	20-29	30-39	40-49
Gastric cancer			
Random	648/0 =	2591/3 = 864	6544/3 = 2182
Iron deficiency			
Yes	379/0 =	1656/3 = 552	4592/1 = 4592
No	269/0 =	935/0 =	1952/2 = 976
Any symptom			
Yes	319/0 =	1313/3 = 438	3446/3 = 1149
No	329/0 =	1278/0 =	3098/0 =
Weight loss			
Yes	23/0 =	53/1 = 53	144/2 = 72
No	625/0 =	2538/2 = 1269	6400/1 = 6400
Any gastric symptoms: UGI and/or dysphagia			
<i>All women</i>			
Yes	214/0 =	959/3 = 320	2620/2 = 1310
No	434/0 =	1632/0 =	3924/1 = 3924
<i>Iron deficient (+) women (n = 7919)</i>			
Yes	125/0 =	615/3 = 205	1823/1 = 1823
No	254/0 =	1041/0 =	2769/0 =
<i>Iron deficient (-) women (n = 4165)</i>			
Yes	89/0 =	344/0 =	797/1 = 797
No	180/0 =	591/0 =	1155/1 = 1155
Colon cancer			
Random	648/1 = 648	2591/7 = 371	6544/18 = 364
Iron deficiency			
Yes	379/1 = 379	1656/5 = 332	4592/12 = 383
No	269/0 =	935/2 = 468	1952/6 = 326
Any symptoms			
Yes	319/1 = 319	1313/7 = 188	3446/13 = 266
No	329/0 =	1278/0 =	3098/5 = 620
Weight loss			
Yes	23/0 =	53/0 =	144/1 = 144
No	625/1 = 625	2538/7 = 363	6400/17 = 377
Any colon symptoms: LGI and/or rectal bleed			
<i>All women</i>			

Sample size/n of cancers^a = number needed to detect one cancer	20-29	30-39	40-49
Yes	202/1 = 202	750/7 = 108	1841/11 = 168
No	446/0 =	1841/0 =	4703/7 = 672
<i>Iron deficient (+) women (n = 7919)</i>			
Yes	120/1 = 120	485/5 = 97	1244/8 = 156
No	259/0 =	1171/0 =	3348/4 = 837
<i>Iron deficient (-) women (n = 4165)</i>			
Yes	82/0 =	265/2 = 133	597/3 = 199
No	187/0 =	670/0 =	1355/3 = 452

IDA = iron deficiency anemia, UGI = upper gastrointestinal, LGI = lower gastrointestinal

^a Within five years of the iron deficiency test date.

eTable 6. Unadjusted incidence of esophagitis, ulcers and IBD in women who underwent endoscopy procedure within one year of testing for iron deficiency

	Iron Deficient			Non-Iron Deficient			<i>P</i> Value
	Sample Size (N)	Pathology ^a (n)	Incidence/1000 Patients (95% CI)	Sample Size (N)	Pathology ^a (n)	Incidence/1000 Patients (95% CI)	
Anemic women (N = 6536)							
Esophagitis							
Age 20-49, years	527	39	74.00 (54.61-99.57)	126	12	95.24 (55.32-159.10)	.42
20-29	11	2	181.80 (51.37-477.00)	4	0	--	>.99
30-39	82	10	122.00 (67.61-210.10)	36	7	194.40 (97.53-350.30)	.30
40-49	434	27	62.21 (43.10-89.00)	86	5	58.14 (25.09-129.00)	.89
Ulcer							
Age 20-49, years	527	20	37.95 (24.70-57.89)	126	10	79.37 (43.68-139.90)	.046
20-29	11	0	--	4	0	--	--
30-39	82	6	73.17 (33.96-150.60)	36	2	55.56 (15.37-181.50)	>.99
40-49	434	14	32.26 (19.31-53.41)	86	8	93.02 (47.89-173.00)	.02
IBD							
Age 20-49, years	527	4	7.59 (2.96-19.35)	126	3	23.81 (8.13-67.67)	.14
20-29	11	0	--	4	0	--	
30-39	82	0	--	36	1	27.78 (4.92-141.70)	.31
40-49	434	4	9.22 (3.59-23.46)	86	2	23.26 (6.40-80.88)	.26
Non-Anemic women							

	Iron Deficient			Non-Iron Deficient			P Value
	Sample Size (N)	Pathology ^a (n)	Incidence/1000 Patients (95% CI)	Sample Size (N)	Pathology ^a (n)	Incidence/1000 Patients (95% CI)	
(N = 939)							
Esophagitis							
Age 20-49, years	388	19	48.97 (31.57-75.21)	551	43	78.04 (58.45-103.50)	.08
20-29	33	1	30.30 (5.37-153.20)	47	4	85.11 (33.59-199.30)	.40
30-39	75	4	53.33 (20.93-129.30)	132	7	53.03 (25.92-105.40)	>.99
40-49	280	14	50.00 (30.01-82.17)	372	32	86.02 (61.59-118.90)	.08
Ulcer							
Age 20-49, years	388	19	48.97 (31.57-75.21)	551	24	43.56 (29.44-63.99)	.70
20-29	33	1	30.30 (5.37-153.20)	47	2	42.55 (11.75-142.50)	>.99
30-39	75	4	53.33 (20.93-129.30)	132	6	45.45 (21.00-95.62)	>.99
40-49	280	14	50.00 (30.01-82.17)	372	16	43.01 (26.65-68.72)	.67
IBD							
Age 20-49, years	388	9	23.20 (12.25-43.49)	551	13	23.59 (13.84-39.94)	.97
20-29	33	0	--	47	2	42.55 (11.75-142.50)	.51
30-39	75	2	13.33 (2.36-71.73)	132	2	15.15 (4.16-53.56)	>.99
40-49	280	8	28.57 (14.55-55.36)	372	9	24.19 (12.78-45.33)	.73

IBD= inflammatory bowel disease

- Not applicable

^a Within five years of the iron deficiency test date.

eTable 7. Demographic and clinical characteristics in women age 20-49 without anemia who were tested for iron deficiency.

Characteristics	Iron Deficiency		Had Endoscopy ^a	
	Yes	No	Yes	No
Women age 20-49 (N = 22,657 [100])	n = 6979	n = 15,678	n = 939	n = 21,718
Iron deficiency	--	--	388 (41.3)	6591 (30.4)
Had endoscopy ^a	388 (5.6)	551 (3.5)	--	--
Age groups (year)				
20-29	1010 (14.5)	2122 (13.5)	80 (8.5)	3052 (14.1)
30-39	2068 (29.6)	5076 (32.4)	202 (22.0)	6937 (31.9)
40-49	3901 (55.9)	8480 (54.1)	652 (69.4)	11,729 (54.0)
Age (year), mean \pm SD	38.9 \pm 7.9	39.0 \pm 7.7	41.4 \pm 7.2	38.8 \pm 7.8
Charlson comorbidity index score				
Mean \pm SD	0.4 \pm 0.8	0.5 \pm 1.0	0.8 \pm 1.2	0.5 \pm 0.9
≥ 1	2372 (34.0)	5641 (36.0)	441 (47.0)	7572 (34.9)
Race/Ethnicity				
White	3075 (44.1)	7324 (46.7)	481 (51.2)	9918 (45.7)
Black	520 (7.5)	1382 (8.8)	60 (6.4)	1842 (8.5)
Hispanic	1672 (24.0)	2987 (19.1)	199 (21.2)	4460 (20.5)
Asian	1192 (17.1)	2771 (17.5)	125 (13.3)	3838 (17.7)
Other ^b	520 (7.5)	1214 (7.7)	74 (7.9)	1660 (7.6)
Any comorbidity	1454 (20.8)	2816 (18.0)	211 (22.5)	4059 (18.7)
Diabetes mellitus	558 (8.0)	1370 (8.7)	97 (10.3)	1831 (8.4)
Kidney disease	41 (0.6)	206 (1.3)	17 (1.8)	230 (1.1)
Cardiovascular disease	58 (0.8)	136 (0.9)	10 (1.1)	184 (0.9)
Pulmonary disease	75 (1.1)	388 (2.5)	17 (1.8)	446 (2.1)
Menorrhagia	854 (12.2)	1046 (6.7)	97 (10.3)	1803 (8.3)
Cirrhosis	8 (0.1)	39 (0.3)	9 (1.0)	38 (0.2)
Any symptoms	3810 (54.6)	8504 (54.3)	837 (89.1)	11,477 (52.9)

Characteristics	Iron Deficiency		Had Endoscopy ^a	
	Yes	No	Yes	No
UGI symptoms	2769 (39.7)	6052 (38.6)	659 (70.2)	8162 (37.6)
Dysphagia	187 (2.7)	433 (2.8)	83 (8.8)	537 (2.5)
LGI symptoms	2016 (28.9)	4628 (29.5)	528 (56.2)	6116 (28.2)
Rectal bleed	463 (6.6)	1023 (6.5)	195 (20.8)	1291 (5.9)
Weight loss	160 (2.3)	374 (2.4)	74 (7.9)	460 (2.1)
Any gastric symptoms: UGI and/or dysphagia	2870 (41.1)	6304 (40.2)	684 (72.8)	8490 (39.1)
Any colon symptoms: LGI and/or rectal bleed	2314 (33.2)	5303 (33.8)	630 (67.1)	6987 (32.2)
Endoscopy within 5 years of study entry	1028 (14.7)	1766 (11.3)	939 (100)	1855 (8.5)

Data are column % unless otherwise stated.

-- = Not applicable; SD = standard deviation; UGI = upper gastrointestinal; LGI = lower gastrointestinal.

^a Defined as undergoing endoscopy within one year of the iron deficiency test date.

^b Including American Indians, Alaskan Natives, more than one racial groups, and unknown.

eTable 8. Unadjusted incidence of gastric and colon cancers stratified by iron deficiency and symptoms in 22,657 non-anemic women age 20-49.

	Sample Size (N)	Cancer ^a (n)	NND One Cancer (95% CI)	Incidence/1000 Patients (95% CI)	P Value
Gastric cancer					
Random	22,657	3	7553 (2569-22,207)	0.13 (0.05-0.39)	--
Iron deficiency					
Yes	6979	0	--	--	.56
No	15,678	3	5226 (1778-15,367)	0.19 (0.07-0.56)	
Any symptoms					
Yes	12,314	3	4105 (1397-12,069)	0.24 (0.08-0.72)	.26
No	10,343	3	--	--	
UGI symptoms					
Yes	8821	3	2941 (1001-8646)	0.34 (0.12-1.00)	.06
No	13,836	0	--	--	
Dysphagia					
Yes	620	1	620 (111-3512)	1.61 (0.28-9.08)	.08
No	22,037	2	11,019 (3023-40,179)	0.09 (0.02-0.33)	
LGI symptoms					
Yes	6644	1	6644 (1174-37,638)	--	>.99
No	16,013	2	8007 (2197-29,196)	0.12 (0.03-0.46)	
Rectal bleed					
Yes	1486	1	1486 (264-8418)	0.67 (0.12-3.80)	.18
No	21,171	2	10,586 (2904-38,600)	0.09 (0.03-0.34)	
Weight loss					
Yes	534	1	534 (95-3025)	1.87 (0.33-10.5)	.07
No	22,123	2	11,062 (3035-40,336)	0.09 (0.02-0.33)	
Any gastric symptoms: UGI and/or dysphagia					
All women					
Yes	8968	3	2990 (1018-8790)	0.33 (0.11-0.98)	.06
No	13,689	0	--	--	
Iron deficient (+) women (n = 6979)					
Yes	2801	0	--	--	
No	4178	0	--	--	
Iron deficient (-) women (n = 15,678)					
Yes	6167	3	2056 (700-6044)	0.49 (0.17-1.43)	.06
No	9511	0	--	--	
Colon cancer					
Random	22,657	28	810 (561-1170)	1.24 (0.86-1.79)	--
Iron deficiency					
(+)	6979	16	437 (269-709)	2.29 (1.41-3.72)	<.01
(-)	15,678	12	1307 (748-2284)	0.77 (0.44-1.34)	

	Sample Size (N)	Cancer ^a (n)	NND One Cancer (95% CI)	Incidence/1000 Patients (95% CI)	P Value
Any symptoms					
Yes	12,314	25	493 (334-727)	2.03 (1.38-3.00)	<.01
No	10,343	3	3448 (1174-10,138)	0.29 (0.10-0.85)	
UGI symptoms					
Yes	8821	9	981 (516-1863)	1.02 (0.54-1.94)	.46
No	13,836	19	729 (467-1138)	1.37 (0.88-2.14)	
Dysphagia					
Yes	620	1	620 (111-3512)	1.61 (0.28-9.08)	.54
No	22,037	27	817 (562-1188)	1.23 (0.84-1.78)	
LGI symptoms					
Yes	6644	15	443 (269-731)	2.26 (1.37-3.72)	<.01
No	16,013	13	1232 (721-2108)	0.81 (0.47-1.39)	
Rectal bleed					
Yes	1486	16	93 (58-151)	10.8 (6.64-17.4)	<.01
No	21,171	12	1765 (1010-3084)	0.57 (0.32-0.99)	
Weight loss					
Yes	534	7	77 (38-158)	12.1 (6.36-26.8)	<.01
No	22,123	21	1054 (690-1611)	0.95 (0.62-1.45)	
Any colon symptoms: LGI and/or rectal bleed					
All women					
Yes	7364	19	388 (249-606)	2.58 (1.65-4.03)	<.01
No	15,293	9	1700 (895-3230)	0.59 (0.31-1.12)	
Iron deficient (+) women (n = 6979)					
Yes	2232	10	224 (122-411)	4.48 (2.44-8.23)	.01
No	4747	6	792 (363-1726)	1.26 (0.58-2.76)	
Iron deficient (-) women (n = 15,678)					
Yes	5132	9	571 (301-1084)	1.75 (0.92-3.33)	<.01
No	10,546	3	3516 (1197-10,336)	0.28 (0.10-0.84)	

Comparisons were made using Fisher's exact tests.

NND = number needed to detect; -- = not applicable; (+) = positive; (-) = negative; - = not applicable; UGI = upper gastrointestinal; LGI = lower gastrointestinal.

^a Within five years of the iron deficiency test date.