

Association between socioeconomic status and mucosal healing in Japanese patients with ulcerative colitis: a cross-sectional study

Shogo Kitahata,¹ Shinya Furukawa ,² Teruki Miyake,³ Kana Shiraishi,³ Kazuhiro Tange,⁴ Yu Hashimoto,³ Sen Yagi,⁵ Tomoyuki Ninomiya,¹ Seiyuu Suzuki,⁶ Naozumi Shibata,⁷ Hidehiro Murakami,⁸ Katsuhisa Ohashi,⁹ Hideomi Tomida,¹⁰ Yasunori Yamamoto,¹⁰ Eiji Takeshita,⁴ Yoshio Ikeda,¹⁰ Yoichi Hiasa³

To cite: Kitahata S, Furukawa S, Miyake T, *et al*. Association between socioeconomic status and mucosal healing in Japanese patients with ulcerative colitis: a cross-sectional study. *BMJ Open Gastro* 2022;**9**:e001000. doi:10.1136/bmjgast-2022-001000

► Additional supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/bmjgast-2022-001000>).

SK and SF contributed equally.

SK and SF are joint first authors.

Received 30 July 2022
Accepted 4 October 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Professor Shinya Furukawa;
shinya.furukawa@gmail.com

ABSTRACT

Objective Socioeconomic status is a risk factor for worse outcomes in many diseases. However, evidence on the association between socioeconomic status and clinical outcome in patients with ulcerative colitis (UC) is limited. In the clinical setting, the therapeutic goal for UC is to achieve mucosal healing (MH). Thus, the aim of this study is to examine the association between socioeconomic status and MH in patients with UC.

Methods The study population consisted of 298 patients with UC. Education status and household income were divided into three groups based on a self-administered questionnaire. MH and complete MH were defined as a Mayo endoscopic subscore of 0–1 and 0, respectively. The association of socioeconomic status with MH and complete MH was assessed by multivariate logistic regression analysis. Patients with UC were divided into a younger group (<51 years old) and an older group (≥51 years old) based on median age.

Results The percentage of MH and complete MH was 62.4% and 25.2%, respectively. In all patients, socioeconomic status was not associated with MH and complete MH, respectively. In the older group, education but not household income was independently positively associated with MH and complete MH. In contrast, in the younger group, no association between socioeconomic status and MH and complete MH was found.

Conclusion In older Japanese patients with UC, education status but not household income was independently positively associated with MH and complete MH.

INTRODUCTION

Ulcerative colitis (UC) is a chronic inflammatory bowel disease (IBD) that usually requires long-term, lifelong treatment. The estimated prevalence of UC is increasing in Asian countries and remains high in Western countries.¹ Repeated relapses of UC cause poor quality of life, surgery and the risk of colorectal cancer.² Both cohort studies and randomised controlled trial showed that mucosal healing (MH) is associated with better outcomes.^{2,3}

WHAT IS ALREADY KNOWN ON THIS TOPIC

- ⇒ Several studies of patients with Crohn's disease had revealed an association between socioeconomic status and poor clinical outcomes.
- ⇒ Evidence regarding this issue in patients with ulcerative colitis (UC) is limited.
- ⇒ Additionally, the association regarding this issue in patients with UC is inconsistent.
- ⇒ No study has investigated this issue in Asian patients with UC.

WHAT THIS STUDY ADDS

- ⇒ Higher education was independently positively associated with mucosal healing and complete mucosal healing in older but not younger patients with UC.
- ⇒ In contrast, no association between household income and clinical outcome was found, regardless of age.
- ⇒ This is the first study to show a positive association between education status and clinical outcomes in Asian patients with UC.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

- ⇒ Even Asian patients with UC, high education might be protective factor for poor prognosis in patients with UC.
- ⇒ In older and less educated patients with UC, endoscopies should be performed more frequently, the dose of drug administered should be started higher, and the dose reduction should be slow.

Maintaining MH is associated with a reduced risk of relapse, fewer surgeries, fewer hospitalisation and steroid tapering. Thus, MH is recommended as the therapeutic goal in clinical setting.^{2,3} IBD imposes considerable costs on individuals by affecting their daily lives and social environment. Therefore, socioeconomic status is considered an important

**Table 1** Clinical characteristics of 298 study participants

Variable	n (%)
Age, years, mean±SD	50.8±15.9
Men (%)	174 (58.4)
Disease extent (pancolitis/left-sided/proctitis/ others)	125/80/86/7
Duration of UC, year, mean±SD	8.6±8.7
BMI, kg/m ² , mean±SD	22.81±4.71
Current smoking (%)	25 (8.4)
Current alcohol intake (%)	123 (41.3)
Medication	
5-Aminosalicylates (%)	248 (90.8)
Prednisolone (%)	62 (20.8)
Thiopurines (%)	444 (14.8)
TNF- α monoclonal antibody (%)	18 (6.0)
Mayo endoscopic subscore, mean±SD	1.20±0.91
Mucosal healing (Mayo endoscopic subscore 0 and 1) (%)	186 (62.4)
Complete mucosal healing (Mayo endoscopic subscore 0) (%)	75 (25.2)
Education, years	
Low (\leq 12 years) (%)	149 (50.0)
Moderate (12–16 years) (%)	63 (21.1)
High (\geq 16 years) (%)	86 (28.9)
Household income, Japanese yen/year	
Low (\leq 2 999 999) (%)	63 (21.1)
Moderate (3 000 000–5 999 999) (%)	139 (46.6)
High (\geq 6 000 000) (%)	96 (32.2)
Others: right-sided, segmental colitis and postoperative patients (lack of any preoperative medical records of postoperative patients).	
BMI, body mass index; TNF, tumour necrosis factor; UC, ulcerative colitis.	

factor. Lower socioeconomic status (LSS) might worsen clinical outcomes in many diseases.⁴

Evidence regarding the association between socioeconomic status and clinical outcomes in patients with IBD exists. Most studies of patients with Crohn's disease (CD) showed that education, income and deprivation were associated with poor outcomes.⁵ In a Canadian,⁶ a French,⁷ a British,⁸ two Scottish^{9,10} and three UC studies of patients with CD,^{11–13} socioeconomic status was associated with poor clinical outcomes. In contrast, in patients with UC, evidence regarding the association between socioeconomic status and clinical outcomes is limited and inconsistent. Poor socioeconomic status was associated with mortality in a Canadian study.¹⁴ No association between socioeconomic status and clinical outcomes was found in a British cohort.⁵ In contrast, in a Swedish retrospective study of patients with UC, restorative surgery was more common in patients with high income.¹⁵ The association

between socioeconomic status and clinical outcomes in patients with UC remains inconsistent. Additionally, no study has investigated this issue in Asian patients with UC.

In Japan, education status differs for individuals over 50 years of age due to the influence of economic growth.¹⁶ We hypothesised that education status would affect clinical outcomes in patients with UC and that the association between education and clinical outcomes differs between older and younger patients. In the clinical setting, the therapeutic goal for UC is to achieve MH.^{2,3,17,18} The purpose of this study was to investigate the association between socioeconomic status and MH in Japanese patients with UC.

METHODS

Study design

This was a cross-sectional study using baseline data from a prospective cohort study.

Study population

The subjects were 387 patients with UC who visited the Department of Gastroenterology and Metabology at the Ehime University Graduate School of Medicine or several affiliated hospitals in Ehime Prefecture, Japan, as inpatients or outpatients between 2015 and 2019. All patients consented to this study and were able to answer the self-administered questionnaire. Subjects were diagnosed with UC based on endoscopic findings and met clinical, radiological and histological criteria. The protocol for this study was developed in accordance with the 1964 Declaration of Helsinki and subsequent versions of the ethical guidelines. Well-trained staff obtained written informed consent from all registered patients.

Measurements

A self-administered questionnaire elicited information regarding smoking, drinking, education and household income. Information on medication for UC, disease extent and duration of UC was obtained from patient medical records. Current smoking was deemed positive if patients answered that they were currently smoking, regardless of the number of cigarettes. Current alcohol intake was deemed positive if patients answered that they had a drinking habit, regardless of the frequency of drinking or the amount of alcohol.

Definition of MH and complete MH

In this study, MH and complete MH were defined as Mayo endoscopic score (MES) category 0–1 and 0, respectively.¹⁹ A certified endoscopist evaluated endoscopic activity by total colonoscopy and reported endoscopic findings and key images. Endoscopist regularly participated in meeting to review endoscopic findings. A single endoscope specialist was responsible for evaluating MES, MH and complete MH at almost the same time as this survey and was blinded to other findings including socioeconomic status.

Table 2 Crude and adjusted ORs and 95% CIs for education status and household income relation to mucosal healing

Variable	Prevalence (%)	Crude OR (95% CI)	Adjusted OR (95% CI)
<i>MH (MES 0–1)</i>			
Education, years			
Low (≤ 12 years)	90/149 (60.4)	1.00	1.00
Moderate (12–16 years)	40/63 (63.5)	1.14 (0.62 to 2.12)	1.31 (0.66 to 2.67)
High (≥ 16 years)	56/86 (65.1)	1.22 (0.71 to 2.14)	1.43 (0.77 to 2.69)
P for trend			0.28
Household income, Japanese yen/year			
Low ($\leq 2\,999\,999$)	38/63 (60.3)	1.00	1.00
Moderate (3 000 000–5 999 999)	89/139 (64.0)	1.17 (0.63 to 2.16)	1.39 (0.70 to 2.74)
High ($\geq 6\,000\,000$)	59/96 (61.5)	1.05 (0.55 to 2.09)	1.31 (0.61 to 2.80)
P for trend			0.54
<i>Complete MH (MES 0)</i>			
Education, years			
Low (≤ 12 years)	35/149 (23.5)	1.00	1.00
Moderate (12–16 years)	16/63 (25.4)	1.11 (0.55 to 2.17)	1.63 (0.74 to 3.55)
High (≥ 16 years)	24/86 (27.9)	1.26 (0.68 to 2.30)	1.67 (0.86 to 3.27)
P for trend			0.14
Household income, Japanese yen/year			
Low ($\leq 2\,999\,999$)	20/63 (31.8)	1.00	1.00
Moderate (3 000 000–5 999 999)	38/139 (27.3)	0.81 (0.43 to 1.56)	0.86 (0.42 to 1.76)
High ($\geq 6\,000\,000$)	17/96 (17.7)	0.46 (0.22 to 0.97)	0.47 (0.20 to 1.09)
P for trend			0.07
Adjusted for age, sex, use of prednisolone, current alcohol intake, current smoking, BMI, duration of disease, disease extent (pancolitis/non-pancolitis), education and household income. BMI, body mass index; MES, Mayo endoscopic subscore; MH, mucosal healing.			

Statistical analysis

Education status was classified into three categories: (1) low education (junior high school and high school), ≤ 12 years, (2) moderate education (junior college, vocational technical school and 4-year college dropout), 12–16 years and (3) high education: college or graduate school, 16 years or more. Household income was classified into three categories: (1) low income, $\leq 2\,999\,999$ Japanese yen/year, (2) moderate income, 3 000 000–5 999 999 Japanese yen/year and (3) high income $\geq 6\,000\,000$ Japanese yen/year. Age was divided into younger (< 51 years old) and older (≥ 51 years old) groups based on the mean age (50.8 years). Estimations of crude ORs and their 95% CIs for MH and clinical remission in relation to education were performed using logistic regression analysis. Multiple logistic regression analyses were used to adjust for potential confounding factors. Age, sex, use of prednisolone, current alcohol intake, current smoking, body mass index (BMI), duration of disease, disease extent (pancolitis/non-pancolitis), education and household income were selected a priori as potential confounding factors. All statistical analyses were performed using SAS software package V.9.4 (SAS Institute, Cary, North Carolina, USA). All probability values for statistical tests

were two-tailed, and $p < 0.05$ was considered statistically significant.

RESULTS

The characteristics of this study

Clinical characteristics of this analysis are shown in table 1. Excluding patients with incomplete data, completed questionnaires were analysed for 77% of patients (298/387). The percentage of MH (MES 0–1) and complete MH (MES 0) was 62.4% and 25.2%, respectively. The education period (years) was 50.0% ($n=149$) for ≤ 12 , 21.1% ($n=63$) for 12–16 and 28.9% ($n=86$) for > 16 years, respectively. The percentage of low income, moderate income and high income was 21.1%, 46.6% and 32.2%, respectively. Age was not associated with household income level in this study (data not shown). The percentage of proportion of high and moderate educational levels decreases in patients with 61 years or older; the proportions of patient with low and high education levels were 33.7% and 32.5%, respectively, among those under 40 years old; 48.7% and 30.3%, respectively, among those aged 41–50 years; 38.0% and 36.0%, respectively, among

**Table 3** Crude and adjusted ORs and 95% CIs for education status and household income relation to mucosal healing in the younger group

Variable	Prevalence (%)	Crude OR (95% CI)	Adjusted OR (95% CI)
<51 years old (n=164)			
MH (MES 0–1)			
Education			
Low (≤12 years)	44/67 (65.7)	1.00	1.00
Moderate (12–16 years)	27/45 (60.0)	0.78 (0.36 to 1.72)	0.98 (0.38 to 2.44)
High (≥16 years)	30/52 (57.7)	0.71 (0.34 to 1.51)	0.79 (0.34 to 1.87)
P for trend			0.54
Household income, Japanese yen/year			
Low (≤2 999 999)	15/24 (62.5)	1.00	1.00
Moderate (3 000 000–5 999 999)	51/78 (65.4)	1.13 (0.43 to 2.90)	1.25 (0.40 to 3.76)
High (≥6 000 000)	35/62 (56.5)	0.78 (0.29 to 2.02)	1.03 (0.32 to 3.19)
P for trend			0.89
Complete MH (MES 0)			
Education			
Low (≤12 years)	15/67 (22.4)	1.00	1.00
Moderate (12–16 years)	10/45 (22.2)	0.99 (0.39 to 2.44)	1.13 (0.40 to 3.11)
High (≥16 years)	11/52 (21.2)	0.93 (0.38 to 2.23)	0.97 (0.37 to 2.45)
P for trend			0.90
Household income, Japanese yen/year			
Low (≤2 999 999)	7/24 (29.2)	1.00	1.00
Moderate (3 000 000–5 999 999)	20/78 (25.6)	0.84 (0.31 to 2.43)	0.82 (0.27 to 2.59)
High (≥6 000 000)	9/62 (14.5)	0.41 (0.13 to 1.31)	0.43 (0.13 to 1.48)
P for trend			0.12
Adjusted for age, sex, use of prednisolone, current alcohol intake, current smoking, BMI, duration of disease, disease extent (pancolitis/non-pancolitis), education and household income.			
BMI, body mass index; MES, Mayo endoscopic subscore; MH, mucosal healing.			

those aged 51–60 years and 73.0% and 20.2%, respectively, among those aged 61 years or older.

Association between socioeconomic status and endoscopic activity

The association between socioeconomic status and MH in all patients is shown in table 2. Education status was not associated with MH and complete MH. In crude analysis, high income was inversely associated with complete MH (crude OR 0.46, 95% CI 0.22 to 0.97). After adjustment for confounding factors, the association between high income and complete MH disappeared.

Association between socioeconomic status and endoscopic activity after age stratification at time of survey

Crude and adjusted ORs and 95% CIs for socioeconomic status related to MH and complete MH in the younger group (table 3) and the older group (table 4) are shown. No association between socioeconomic status and clinical outcome was found in the younger group. In the older group, higher education was significantly and positively associated with MH (MH: crude OR 2.54,

95% CI 1.06 to 6.62). After adjustment for age, sex, use of prednisolone, current alcohol intake, current smoking, BMI, duration of disease, disease extent, education and household income, higher education was independently positively associated with MH and complete MH, respectively (MH: adjusted OR 3.27, 95% CI 1.14 to 10.48 and complete MH: adjusted OR 3.19, 95% CI 1.13 to 9.44). After excluding patients with UC diagnosis of 1 year or less, the positive association between education status and MH was significant (data not shown). In contrast, no association between household income and clinical outcome was found.

DISCUSSION

In this study, higher education was independently positively associated with MH and complete MH in older but not younger patients with UC. In contrast, no association between household income and clinical outcome was found, regardless of age. This is the first study to show a positive association between education status and clinical outcomes in Asian patients with UC.

Table 4 Crude and adjusted ORs and 95% CIs for education status and household income relation to mucosal healing in the older group

Variable	Prevalence (%)	Crude OR (95% CI)	Adjusted OR (95% CI)
≥51 years old (n=134)			
<i>MH (MES 0–1)</i>			
Low (≤12 years)	46/82 (56.1)	1.00	1.00
Moderate (12–16 years)	13/18 (72.2)	2.04 (0.70 to 6.82)	2.67 (0.76 to 10.64)
High (≥16 years)	26/34 (76.5)	2.54 (1.06 to 6.62)	3.27 (1.14 to 10.48)
P for trend			0.020
Household income, Japanese yen/year			
Low (≤2 999 999)	23/39 (59.0)	1.00	1.00
Moderate (3 000 000–5 999 999)	38/61 (62.3)	1.15 (0.50 to 2.62)	1.20 (0.47 to 3.06)
High (≥6 000 000)	24/34 (70.6)	1.67 (0.64 to 4.53)	1.24 (0.37 to 4.29)
P for trend			0.67
<i>Complete MH (MES 0)</i>			
Education			
Low (≤12 years)	20/82 (24.4)	1.00	1.00
Moderate (12–16 years)	6/18 (33.3)	1.55 (0.49 to 4.56)	3.02 (0.77 to 11.81)
High (≥16 years)	13/34 (38.2)	1.92 (0.81 to 4.52)	3.19 (1.13 to 9.44)
P for trend			0.027
Household income, Japanese yen/year			
Low (≤2 999 999)	13/39 (33.3)	1.00	1.00
Moderate (3 000 000–5 999 999)	18/61 (29.5)	0.84 (0.35 to 2.01)	0.81 (0.30 to 2.23)
High (≥6 000 000)	8/34 (23.5)	0.62 (0.21 to 1.71)	0.41 (0.11 to 1.44)
P for trend			0.19
Adjusted for age, sex, use of prednisolone, current alcohol intake, current smoking, BMI, duration of disease, disease extent (pancolitis/non-pancolitis), education and household income.			
BMI, body mass index; MES, Mayo endoscopic subscore; MH, mucosal healing.			

Evidence of an association between socioeconomic status, including household income status, and clinical outcomes in patients with IBD exists (online supplemental file 1). In a Canadian database study of 9298 patients with IBD, LSS was associated with intensive care unit admission, hospital mortality, surgery and high dose steroid use.⁶ In a US study of 323 patients with IBD including 138 patients with UC, low annual income but not high education was positively associated with moderate to severe disability based on IBD Disability Index score.¹³

In most studies of patients with CD, poor socioeconomic status might worsen clinical outcomes. In a French prospective study of 207 patients with CD, deprivation was associated with hospitalisation.⁷ In a US retrospective study of 944 patients with CD, low household income was associated with the hospitalisation rate.¹¹ In a US database study of patients with CD, above-average income was associated with lower in-hospital mortality.¹² In a British study, no association between social deprivation and mortality is found, while the hospitalised prevalence of CD was slightly higher in the most deprived area.⁸ In a Scottish study of 1595 inpatients with CD, social deprivation was significantly associated with 3 years mortality.⁹

Another Scottish database study showed that deprivation status was associated with 3-year mortality after hospitalisation.¹⁰

In contrast, limited evidence regarding socioeconomic status and clinical outcome in patients with UC exists, and their association has been inconsistent. A British study showed that no association between living in socially disadvantaged areas and clinical outcome including hospitalisation and mortality.⁸ In a Canadian population-based study of patients with IBD, LSS was associated with mortality in patients with UC.¹⁴ In a Swedish retrospective study of 5969 patients with UC, no association between socioeconomic status and the risk of failure after restorative surgery was found, while restorative surgery was more common in patients with high income but not education level.¹⁵ The discrepancies between the findings in this study and those of previous studies can be partly explained by age, education status, income status, definition of clinical outcome and medical systems.

The mechanisms underlying the association between education status and clinical outcome remain unclear. LSS affects medication adherence²⁰ and access to health-care.²¹ In a Japanese study of patients with chronic



disease, advanced education is associated with good adherence.²² Similarly, in patients with diabetes, adherence is associated with high education.²³ However, Indian study of patients with IBD, poor adherence was observed in patients with professional or honours and middle school certificate/primary school, while good adherence is found in patients with intermediate or post high school diploma/high school certificate, graduate or post-graduate and illiterate.²⁴ In a French mail-based studies of patients, education is not associated with adherence.²⁵ The association between education status and adherence is inconsistent. In an Indian study, the first reason for non-adherence was reported as forgetting the medication, and the third reason was feeling well. The good clinical course may be making patients less aware of the need for treatment, resulting in lower adherence. A considerable heterogeneity in the adherence rates had been reported in patients with IBD.²⁶ In Japan, the percentage of advanced education in the young population is higher than that in the older population. Therefore, at younger ages, differences in education may have a disappearing influence on clinical outcomes. In Japan, the medical insurance system provides standard medication for all patients with UC, regardless of household income. Additionally, the public medical expense subsidy for UC might mask the association between household income and clinical outcomes in UC. Advanced education might lead to more MH via more access to examinations and healthcare. The mechanism of why education is preventive is unknown, and research on the identification of unknown factors associated with education is needed.

In several studies of patients with CD, low socioeconomic status is associated with poor clinical outcomes. In the contrast, in studies of patients with UC, the association between socioeconomic status and clinical outcomes is inconsistent. Socioeconomic status might affect the clinical outcomes in CD more than those in UC.⁵ However, few studies on this topic were conducted in studies of patients with UC. In Japan, the universal health insurance system and subsidies for UC treatment have greatly reduced the high cost of UC treatment, and hospitalisation and surgery have little impact on income. Japan's universal health insurance system and social security might weaken the influence of socioeconomic status to clinical outcomes. Additionally, social health insurance differs from country to country, further more studies regarding this issue in patients with UC is needed.

This study has some limitations. First, the cross-sectional study design failed to demonstrate a causal relationship between education status and UC disease activity. Second, this study used a self-administered questionnaire. Although there are no data on the validity of self-reports of education status and household income, any possible misclassification of non-discriminatory exposure would introduce a bias toward the null hypothesis. We could not assess the reduction of household income due to hospitalisation and/or intensified treatment. Low socioeconomic status may worsen disease activity.

Hospitalisation and absence from work might worsen the socioeconomic status. However, in Japan, the universal health insurance system and subsidies for UC treatment have greatly reduced the high cost of UC treatment, and hospitalisation and surgery have little impact on income. Japan's universal health insurance system and social security might mask the true association between household income and clinical outcomes. Third, although employment status is strongly associated with socioeconomic status, data regarding patient employment status is lacking. Fourth, education, income and medical systems differ among countries. Thus, generalisation of the findings of this study might be difficult for other ethnic populations. Fifth, we did not use central reading of endoscopic findings in the study. Sixth, we could not assess other clinical outcomes including hospitalisation, operation and mortality due to sample size. Finally, selection bias may have influenced the results of this study. The percentage of biologics use might be low in this cohort. This study was probably not representative of all Japanese patients with UC. However, in Japan, medical checkups are widely conducted to screen for colorectal cancer screening using faecal occult blood test. Endoscopic examination is performed for subjects with faecal occult blood test positive, regardless of symptoms. The cohort may include patients with UC with no symptoms but only positive faecal occult blood. Japanese medical checkups might detect no symptomatic patients with UC and lower the percentage of biologics use in the cohort. However, the mean age, sex ratio and drug dosage in this study were similar to those in Japanese national surveys on UC.²⁷

In conclusion, education status but not household income was independently positively associated with MH and complete MH in older Japanese patients with UC. In younger Japanese patients with UC, socioeconomic status might not be associated with clinical outcome. In older and less educated patients with UC, endoscopies should be performed more frequently, the dose of drug administered should be started higher, and the dose reduction should be slow.

Author affiliations

¹Department of Gastroenterology, Ehime Prefectural Central Hospital, Matsuyama, Japan

²Health Services Center, Ehime University, Matsuyama, Japan

³Department of Gastroenterology and Metabolism, Ehime University Graduate School of Medicine, Toon, Japan

⁴Department of Inflammatory Bowel Diseases and Therapeutics, Ehime University Graduate School of Medicine, Toon, Japan

⁵Department of Internal Medicine, Saiseikai Imabari Hospital, Imabari, Japan

⁶Department of Gastroenterology, Medical Corporation Sumitomo Besshi Hospital, Niihama, Japan

⁷Department of Gastroenterology, Ehime Prefectural Niihama Hospital, Niihama, Japan

⁸Department of Internal Medicine, Saiseikai Matsuyama Hospital, Matsuyama, Japan

⁹Ohashi Clinic, Niihama, Japan

¹⁰Endoscopy Center, Ehime University, Matsuyama, Japan

Acknowledgements The authors would like to thank Masakazu Hanayama, Keitarou Kawasaki, Yuji Mizukami, Satoshi Imamine, Masamoto Torisu, Harumi Yano, Makoto Yano, Masato Murakami, Masumi Hino and Tomo Kogama.

Contributors SK, SF and YH contributed to conception and design. KS, KT, YH, SY, TN, SS, NS, HM, KO, HT, YY, ET and YI contributed to material preparation and data collection. SK and SF contributed to data analysis. SK, SF, TM and YH contributed to interpretation of data. The first draft of the manuscript was written by SK and SF. YH contributed to supervision. All authors read and approved the final manuscript. SF was guarantor of this study.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. This study was approved by the ethics committee of Ehime University of Medical Science (approval no. 1505011). Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Shinya Furukawa <http://orcid.org/0000-0002-0041-7688>

REFERENCES

- Ng SC, Shi HY, Hamidi N, *et al.* Worldwide incidence and prevalence of inflammatory bowel disease in the 21st century: a systematic review of population-based studies. *Lancet* 2017;390:2769–78.
- Turner D, Ricciuto A, Lewis A, *et al.* STRIDE-II: an update on the selecting therapeutic targets in inflammatory bowel disease (STRIDE) initiative of the International organization for the study of IBD (IOIBD): determining therapeutic goals for treat-to-target strategies in IBD. *Gastroenterology* 2021;160:1570–83.
- Peyrin-Biroulet L, Sandborn W, Sands BE, *et al.* Selecting therapeutic targets in inflammatory bowel disease (STRIDE): determining therapeutic goals for treat-to-target. *Am J Gastroenterol* 2015;110:1324–38.
- Fiscella K, Franks P, Gold MR, *et al.* Inequality in quality: addressing socioeconomic, racial, and ethnic disparities in health care. *JAMA* 2000;283:2579–84.
- Wardle RA, Wardle AJ, Charadva C, *et al.* Literature review: impacts of socioeconomic status on the risk of inflammatory bowel disease and its outcomes. *Eur J Gastroenterol Hepatol* 2017;29:879–84.
- Bernstein CN, Wald R, Marrie RA. Social determinants of outcomes in inflammatory bowel disease. *Am J Gastroenterol* 2020;115:2036–46.
- Nahon S, Lahmek P, Macaigne G, *et al.* Socioeconomic deprivation does not influence the severity of crohn's disease: results of a prospective multicenter study. *Inflamm Bowel Dis* 2009;15:594–8.
- Button LA, Roberts SE, Goldacre MJ, *et al.* Hospitalized prevalence and 5-year mortality for IBD: record linkage study. *World J Gastroenterol* 2010;16:431–8.
- Kennedy NA, Clark DN, Bauer J, *et al.* Nationwide linkage analysis in Scotland to assess mortality following hospital admission for crohn's disease: 1998–2000. *Aliment Pharmacol Ther* 2012;35:142–53.
- Ventham NT, Kennedy NA, Duffy A, *et al.* Nationwide linkage analysis in Scotland—has mortality following hospital admission for crohn's disease changed in the early 21st century? *J Crohns Colitis* 2014. doi:10.1016/j.crohns.2014.09.002. [Epub ahead of print: 26 Sep 2014].
- Walker C, Allamneni C, Orr J, *et al.* Socioeconomic status and race are both independently associated with increased hospitalization rate among crohn's disease patients. *Sci Rep* 2018;8:4028.
- Nguyen GC, Bayless TM, Powe NR, *et al.* Race and health insurance are predictors of hospitalized crohn's disease patients undergoing bowel resection. *Inflamm Bowel Dis* 2007;13:1408–16.
- Agrawal M, Cohen-Mekelburg S, Kayal M, *et al.* Disability in inflammatory bowel disease patients is associated with race, ethnicity and socio-economic factors. *Aliment Pharmacol Ther* 2019;49:564–71.
- Bernstein CN, Nugent Z, Targownik LE, *et al.* Predictors and risks for death in a population-based study of persons with IBD in manitoba. *Gut* 2015;64:1403–11.
- Nordenvall C, Westberg K, Söderling J, *et al.* Restorative surgery is more common in ulcerative colitis patients with a high income: a population-based study. *Dis Colon Rectum* 2021;64:301–12.
- Ministry of Education, Culture, Sports, Science and Technology of Japan. School basic survey. Available: https://www.mext.go.jp/b_menu/toukei/chousa01/kihon/1267995.htm [Accessed 05 Mar 2022].
- Peyrin-Biroulet L, Ferrante M, Magro F, *et al.* Results from the 2nd scientific workshop of the ECCO. I: impact of mucosal healing on the course of inflammatory bowel disease. *J Crohns Colitis* 2011;5:477–83.
- Shah SC, Colombel J-F, Sands BE, *et al.* Mucosal healing is associated with improved long-term outcomes of patients with ulcerative colitis: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol* 2016;14:1245–55.
- Schroeder KW, Tremaine WJ, Ilstrup DM. Coated oral 5-aminosalicylic acid therapy for mildly to moderately active ulcerative colitis. A randomized study. *N Engl J Med* 1987;317:1625–9.
- Goodhand JR, Kamperidis N, Sirwan B, *et al.* Factors associated with thiopurine non-adherence in patients with inflammatory bowel disease. *Aliment Pharmacol Ther* 2013;38:1097–108.
- Andrulis DP. Access to care is the centerpiece in the elimination of socioeconomic disparities in health. *Ann Intern Med* 1998;129:412–6.
- Ueno H, Yamazaki Y, Yonekura Y, *et al.* Reliability and validity of a 12-Item medication adherence scale for patients with chronic disease in Japan. *BMC Health Serv Res* 2018;18:13–29.
- Kirkman MS, Rowan-Martin MT, Levin R, *et al.* Determinants of adherence to diabetes medications: findings from a large pharmacy claims database. *Diabetes Care* 2015;38:604–9.
- Tomar SK, Kedia S, Singh N, *et al.* Higher education, professional occupation, and upper socioeconomic status are associated with lower adherence to medications in patients with inflammatory bowel disease. *JGH Open* 2019;3:302–9.
- Nahon S, Lahmek P, Saas C, *et al.* Socioeconomic and psychological factors associated with nonadherence to treatment in inflammatory bowel disease patients: results of the ISSEO survey. *Inflamm Bowel Dis* 2011;17:1270–6.
- Vangeli E, Bakhshi S, Baker A, *et al.* A systematic review of factors associated with non-adherence to treatment for immune-mediated inflammatory diseases. *Adv Ther* 2015;32:983–1028.
- Matsuoka K, Igarashi A, Sato N, *et al.* Trends in corticosteroid prescriptions for ulcerative colitis and factors associated with long-term corticosteroid use: analysis using Japanese claims data from 2006 to 2016. *J Crohns Colitis* 2021;15:358–66.

1 Supplement file. Socioeconomic status and clinical outcomes in patients with IBD

Nation	Years	Subjects, number	Socioeconomic status	Clinical outcomes
Canada ⁶	2020	IBD, 9,298	LSS	Physician visits, hospitalization, intensive care unit administration, use of steroid, and death
France ⁷	2009	CD, 207	Deprivation	Hospitalization, and surgery
UK ⁸	2010	CD, 1,467 UC, 1482	Deprivation area	CD, hospitalization UC, no association
Scotland ⁹	2011	Inpatient CD, 1,595	Social Deprivation	3-years mortality
Scotland ¹⁰	2014	CD	Deprivation	Mortality
US ¹¹	2018	CD, 944	Low income, and below poverty level	Hospitalization
US ¹²	2007	Inpatient CD, 41,918	Insurance status, residence area (above median income)	Insurance, low surgery Area, low in-hospital mortality
US ¹³	2019	CD, 185 UC, 138	Household income	Moderate to severe disability
Canada ¹⁴	2015	CD and UC, 10,788	LSS	Mortality
Sweden ¹⁵	2021	UC patients with colectomy, 5,969	High income and education	Income, high rete restorative surgery (No association, fail of surgery) Education, no association

2 LSS, lower socioeconomic status; CD, Crohn's disease; UC, ulcerative colitis