

Supplementary Table 2

	Patients with Abnormal ALT (>1x ULN) (n=152)	Patients with all normal Liver enzymes (n=788)	P-value
Age	56.7 ± 11.9	59.4 ± 13.6	0.02
Sex			<0.01
M	101 (66.5%)	329 (41.8%)	
F	51 (33.6%)	459 (58.3%)	
BMI	30.7 ± 6.2	28.1 ± 5.6	<0.01
Obesity (BMI≥30)	73 (48.3%)	235 (29.9%)	<0.01
Alcohol intake (Within limits: ≤21 units M, ≤14 units F)			<0.01
Non-drinker	23 (15.3%)	164 (21.4%)	
Within limits	98 (65.3%)	528 (68.8%)	
Above limits	29 (19.3%)	75 (9.8%)	
AUDIT – C			<0.01
Low risk	61 (40.4%)	480 (61.3%)	
High risk	90 (59.6%)	302 (38.6%)	
Hypertension	58 (38.2%)	254 (32.3%)	0.16
Type II Diabetes	24 (15.8%)	93 (11.8%)	0.17
Metabolic syndrome (Harmonised criteria)	81 (55.5%)	312 (40.3%)	<0.01
Triglyceride	1.8 ± 1.3	1.2 ± 0.7	<0.01
HDL-cholesterol	1.3 ± 0.4	1.5 ± 0.5	<0.01
FLI	74.4 ± 27.5	47.3 ± 30.2	<0.01
FLI group			<0.01
Low	18 (12.7%)	283 (36.6%)	
Intermediate	16 (11.3%)	208 (26.9%)	

High	108 (76.1%)	283 (36.6%)	
FIB-4 score	1.54 ± 1.04	1.27 ± 0.64	<0.01
Age-adjusted FIB-4 score			<0.01
Low	74 (52.5%)	618 (79.1%)	
Intermediate	57 (40.4%)	139 (17.8%)	
High	10 (7.1%)	24 (3.1%)	
QRisk2	15.8 ± 14.0	15.70 ± 12.8	0.95

Supplementary Table 1: Comparison of patient demographics between patients with abnormal ALT and patients with all normal liver enzymes